DOI: 10.1111/1365-2435.13577

RESEARCH ARTICLE



Impacts of global environmental change drivers on non-structural carbohydrates in terrestrial plants

Ying Du 💿 | Ruiling Lu 💿 | Jianyang Xia 回

Zhejiang Tiantong Forest Ecosystem National Observation and Research Station. Shanghai Key Lab for Urban Ecological Processes and Eco-Restoration, Research Center for Global Change and Ecological Forecasting, School of Ecological and Environmental Sciences, East China Normal University, Shanghai, China

Correspondence

Jianyang Xia Email: jyxia@des.ecnu.edu.cn

Funding information

National Natural Science Foundation of China, Grant/Award Number: 31722009 and 41630528; National 1000 Young Talents Program of China

Handling Editor: Adam Martin

Abstract

- 1. Non-structural carbohydrates (NSCs, including soluble sugars and starch) are essential to support the growth and survival of terrestrial plants. Starch and sugars play different roles in multiple plant ecological functions such as drought tolerance, growth and plant defence, and several other processes which are being rapidly shaped by global environmental change. However, it is uncertain whether soluble sugars and starch show different responses across plant functional types, tissue types and treatment conditions (i.e. the intensity and duration of environmental variability) to global-change drivers.
- 2. Here based on a database of 275 plants (including 17 plant functional types), we conducted a meta-analysis to examine the effects of elevated atmospheric CO₂ concentration (eCO₂), nitrogen (N) addition, drought and warming on NSCs and its components.
- 3. We found NSCs responses to global environmental change were mainly driven by (a) soluble sugar changes in response to N addition and drought, as well as (b) starch changes in response to eCO₂ and warming. The different responses between soluble sugars and starch were more evident under eCO₂ and drought, especially in herbs or leaves. Interactive effects of multiple environmental change drivers on soluble sugars and starch were mainly additive. The divergent main and interactive effects on soluble sugars and starch depend on experimental conditions. For example, the starch responses to eCO₂ and its interaction with N addition were the strongest in short-term experiments.
- 4. Overall, our study shows the divergent responses of soluble sugars and starch in terrestrial plants to different global environmental change drivers, suggesting a changed carbon sink-source balance in plants under future global changes. The findings also highlight that predicting plant functional changes into the future requires a mechanistic understanding of how NSCs and its components are linked to specific environmental change drivers.

KEYWORDS

CO₂, drought, nitrogen addition, non-structural carbohydrates, soluble sugars, starch, warming

1 | INTRODUCTION

Non-structural carbohydrates (NSCs) are crucial to fulfilling multiple functions in plants, including metabolism, transport, osmoregulation and export as substrates for soil organisms (Hartmann & Trumbore, 2016; Smith & Smith, 2011). NSCs are largely comprised of soluble sugars and starch, which have distinct ecological functions. Soluble sugars are mainly synthesized from newly assimilated carbon by daytime photosynthesis and translocated to different organs by mass flow (Kozlowski, 1992). They are crucial in metabolism, osmotic regulation, translocation of energy, signal transduction as well as synthesis of defence chemicals, such as monoterpene olefins (turpentine) and diterpene resin acids (rosin) (Chaves, Maroco, & Pereira, 2003; De Schepper, De Swaef, Bauweraerts, & Steppe, 2013; Hartmann & Trumbore, 2016; Plavcová & Jansen, 2015; Rolland, Baena-Gonzalez, & Sheen, 2006; Steele, Lewinsohn, & Croteau, 1995). Soluble sugars are converted to starch when carbon assimilation exceeds carbon demand for growth and maintenance (Chapin III, Schulze, & Mooney, 1990). The starch can be converted back to soluble sugars when carbon assimilation is insufficient to support metabolic activities (e.g. at night; Chapin III et al., 1990; Gibon et al., 2009; Sulpice et al., 2014). Starch is of great importance because its degradation can produce energy, sugars and metabolites to enhance plant resilience under stressful conditions (Thalmann & Santelia, 2017), and its synthesis can reserve carbon for future use and lower the concentration of soluble sugars, avoiding more respiratory consumption (Chapin III et al., 1990). In short, soluble sugars act as a short-term NSCs reserve for immediate functions and starch is mainly a long-term storage reserve (Dietze et al., 2014; Hartmann & Trumbore, 2016; Thalmann & Santelia, 2017).

Global rises in atmospheric CO₂ concentration, nitrogen (N) deposition rate, air temperature and drought occurrence have become the major external drivers for many recent changes in global patterns of plant growth and survival (Hughes, 2000; Parmesan & Yohe, 2003; Reich, Hobbie, & Lee, 2014; Xu et al., 2013). For example, the increasing atmospheric CO₂ concentration and N deposition have driven the enhancing vegetation growth and greenness over global land areas during the past three decades (Huang et al., 2018; Zhu et al., 2016), whereas climate warming and the associated drought have increased the rate of tree mortality at the global scale (McDowell & Allen, 2015). In order to cope with these environmental changes, plants need to develop different strategies to keep the carbon sink-source balance by adjusting the use of NSCs (Dietze et al., 2014; Körner, 2003). Thus, how NSCs and its components respond to these global-change factors is critical in determining the terrestrial carbon feedback to climate changes (Dietze et al., 2014; Hartmann & Trumbore, 2016; Martínez-Vilalta et al., 2016).

In the past decades, many new insights about NSCs' responses to environmental changes have been reported by experimental studies. Those experimental data have been synthesized by a few metaanalyses to explore general response patterns of NSCs to different global environmental change drivers. For example, Li et al. (2018) has synthesized data of 71 trees species and found more enhancement of starch than soluble sugars in leaves under eCO_2 . However, the stimulation induced by eCO_2 is higher in soluble sugars than starch in a meta-analysis of 31 plant species, which include 7 gymnosperms and 24 angiosperms (8 woody and 16 herbaceous species; Zvereva & Kozlov, 2006). Starch is decreased more than soluble sugars in roots of trees under N addition (Li et al., 2018), but the contrary response pattern of a larger reduction in soluble sugars than starch has been reported under warming (Zvereva & Kozlov, 2006). It has been reported that soluble sugars are increased yet starch is decreased under drought in both leaves and roots of trees (Adams et al., 2017; Li et al., 2018).

However, it is uncertain whether the observed pattern is consistent between tissue types, plant functional types and experimental conditions such as treatment intensities and durations. The leaf is the productive organ in plants and has higher metabolic activities than roots and stems (Brüggemann et al., 2011). Sugars are loaded through lignified and suberized walls, which can be developed as a barrier to limit spread of sugars progressively (Jacobsen, Fisher, Maretzki, & Moore, 1992; Rae, Perroux, & Grof, 2005; Slewinski, 2012). Compared with woody plants, herbaceous plants have less lignified tissues. Hence, one can expect that soluble sugars and starch in leaves or herbaceous plants are more sensitive to environmental changes. Plants can guickly acclimate or adapt to global changes such as climate warming (Atkin & Tjoelker, 2003). Plants grown under CO₂ enhancement are limited by other resources in the long-term, such as N (Reich et al., 2014), but the consumption of NSCs is accelerated under long-term severe drought (e.g. McDowell, 2011). Thus, we further hypothesize that the responses of soluble sugars and starch could be stronger under short- than long-term treatment.

Different global change factors can interact with each other to affect NSCs and its two major components. For example, Huttunen, Saravesi, Markkola, and Niemelä (2013) has found that N addition can alleviate the positive effect of elevated CO₂ and the negative impact of warming on NSCs in Betula pendula respectively. Similarly, the combination of elevated CO₂ and temperature can mitigate the main positive effects of CO2 enhancement and the negative effects of warming on starch in Panicum maximum (Habermann et al., 2019). Some other studies have reported that the negative impact of drought on NSCs can be alleviated by CO₂ enrichment or N addition but exacerbated by warming (Agami, Alamri, El-Mageed, Abousekken, & Hashem, 2018; Duan et al., 2013). However, it is uncertain whether these combined effects of global change drivers are additive or not. The additive interaction means the interactive impact of multiple drivers has no significant difference from the sum of individual impacts (Yue, Fornara, Yang, Peng, Peng, et al., 2017). To date, multiple meta-analyses have found that additive effects are dominant in plant C pool, N concentration, phosphorus (P) pool and C:N:P stoichiometry (Yuan & Chen, 2015; Yue, Fornara, Yang, Peng, Li, et al., 2017; Yue, Fornara, Yang, Peng, Peng, et al., 2017; Yue et al., 2018). Also, Li et al. (2018) have found that the interactions of eCO₂ with nitrogen or drought on NSCs and its components in trees are predominantly additive. Hence, we hypothesize

the additive interactions are dominant across various combinations of global-change factors. Also, few studies have explored whether these interactions are consistent across different durations of environmental changes. It limits our understanding of how global environmental changes affect carbon cycle in terrestrial ecosystems.

Here a comprehensive meta-analysis focusing on NSCs' responses to the primary global environmental change drivers, including eCO_2 , N addition, drought and warming, was conducted based on 341 published studies (see Appendix S1). Our study aims to answer three specific questions, including (a) how soluble sugars and starch differently respond to global-change drivers?; (b) how their responses vary with plant functional type, tissue type and experimental conditions; and (c) whether and how different environmental change drivers interact to impact NSCs and its components?

2 | MATERIALS AND METHODS

2.1 | Data compilation

Peer-reviewed journal articles reporting NSCs and its components in response to multiple environmental change drivers (i.e. elevated CO₂, nitrogen addition, drought and warming) were searched in Web of Science and Google Scholar. The search term combinations were as follows: (non-structural carbohydrates or nonstructural carbohydrates or soluble sugar or starch) and (nitrogen or CO₂ or temperature or warming or drought or water) and (plant or production or productivity or biomass or mass). Moreover, we set the following criteria to minimize the potential uncertainties and select proper observations: (a) experimental duration and magnitude were clearly indicated; (b) full-factorial design was used to examine interactive effects of multiple global change factors; (c) the means, standard deviations/ errors and samples sizes of the selected variables in both of the control and experimental groups could be extracted directly from text, tables or figures using GetData Graph Digitizer 2.24 (http://getdata-graphdigitizer.com/). For studies with multiple levels of experimental treatments, multiple measurements in time, multiple plant species and plant parts, they were considered as multiple independent observations. Overall, 341 articles published before March 2019 were compiled into the literature database (see Appendix S1). It should be noted that there are six studies to use plants grown at alpine or boreal system, but they did not impact the response patterns of NSCs and its components to experimental warming.

To further assess the main and interactive effects of multiple environmental change drivers on NSCs and its components, relevant experimental information was complied. In our database, plants were divided into several groups based on growth forms (woody and herbaceous plants, or tree, shrub, grass and forb), life history (annual, biennial and perennial herb; the results of biennial plants were not shown in results because of limited data), photosynthetic pathways (C_3 , C_4 and CAM herbs) and other functional types (broadleaved and coniferous trees, evergreen and deciduous trees or legume and nonlegume). In order to detect responses of NSCs and its components to different treatment intensities, we grouped them into three categories: 'low intensity', 'moderate intensity' and 'high intensity'. For example, according to enhanced magnitude, elevated CO₂ concentration experiments could be divided into low (≤300 p.p.m.), moderate (>300 and ≤600 p.p.m.) and high (>600 p.p.m.) intensity. For N addition experiments, they were classified into low (≤200%), moderate (>200% and ≤400%) and high (>400%) intensities based on comparison of N addition amount between experimental and control groups within a given study (Li et al., 2018). However, for remaining experiments where control groups were not fertilized with nitrogen, they were grouped into low (<50 kg N/ha), moderate (≥50 and <150 kg N/ha) and high (≥150 kg N/ha) intensities. Among studies, experimental drought decreased water by 14%-100%, which could be classified into low (<50%), moderate (≥50% and <70%) and high (\geq 70%) intensity. Warming experiments were grouped as low (\leq 3°C), moderate (>3°C and ≤6°C) and high (>6°C) intensity based on warming magnitude. The studies were conducted in greenhouse, growth chamber, pot, Free-Air CO₂ Enrichment (FACE) and natural habitats. Hence, we grouped them into laboratory conditions (greenhouse, growth chamber, pot) and field conditions (FACE and natural habitats; Xu, Yan, & Xia, 2019).

2.2 | Data analysis

Owing to differences in quantification methods, NSCs cannot be compared directly among laboratories (Landhäusser et al., 2018; Quentin et al., 2015). However, relative changes of NSCs between experimental and control groups are comparable among studies (Adams et al., 2017; Li et al., 2018; Liu et al., 2016). Hence, we calculated the natural logarithm of the response ratio (In RR) and Hedges' *d* to measure the responses of NSCs and its components to individual and interactive environmental change drivers respectively (Hedges, Gurevitch, & Curtis, 1999). The In RR and Hedges' *d* can both reflect the relative changes of variables. For these test statistics, a value larger than zero represents a positive treatment effect on NSCs and its components. The RR is defined as the ratio of the mean of NSCs and its components in the experimental group (X_e) to that in control group (X_c). The log transformation was to improve statistical behaviour (Equation 1):

$$\ln RR = \ln \left(\frac{X_{\rm e}}{X_{\rm c}} \right). \tag{1}$$

The variance (v_1) was estimated by Equation 2:

$$v_1 = \frac{S_e^2}{N_e X_e^2} + \frac{S_c^2}{N_c X_c^2},$$
 (2)

where S_e and S_c are standard deviations of NSCs and its components in the experimental and control group respectively. N_e and N_c represent sample size in the corresponding groups. The reciprocal of the variance (v₁) was considered as the weight (w₁) of each ln RR (Equation 3). A nonparametric weighting function was used to calculate mean effect size because some studies include two or more observations (Hedges et al., 1999; Li et al., 2018; Liu et al., 2016). The w'_1 is the average of the w_1 over the total number of observations (*n*) in a study (Equation 4). Then, a fixed-effect model (Equation 5) was conducted to calculate the mean effect size and generate 95% confidence interval in MetaWin 2.1 (Rosenberg, Adams, & Gurevitch, 2000):

$$w_1 = \frac{1}{v_1},\tag{3}$$

$$w'_1 = \frac{w_1}{n},$$
 (4)

$$\overline{\ln RR} = \frac{\sum_{i} w'_{i} \times \ln RR_{i}}{\sum_{i} w'_{i}},$$
(5)

where w'_i represents the weight of the *i*th study. The main influence on NSCs and its components was evaluated as significant if the 95% confidence intervals (95% Cl) did not overlap with zero. A resampling bootstrapping method based on 9,999 iterations was used if the sample size was lower than 20.

Furthermore, a random-effect model was used to compare heterogeneity of NSCs and its components within (Q_w) and between (Q_b) categories (i.e. different plant functional types, tissue types, experimental intensities and approaches; Borenstein, Hedges, Higgins, & Rothstein, 2009; Li et al., 2018). The random-effect model assumes the effect size is different among studies and considers the variance of each effect size as a sum of within-study variance and between-study variance (Borenstein, Hedges, Higgins, & Rothstein, 2010). A significant between-group heterogeneity $(Q_{\rm h})$ suggested that the effect size differed among categories (Hedges et al., 1999). The meta-regression considered different weights of NSCs and is components (Thompson & Higgins, 2002). It was used to explore the relationship between NSCs and its components as well as the relationship between soluble sugars or starch and experimental duration. The meta-regression was conducted with the METAFOR package in R 3.4 (R Development Core Team, 2016). The difference in regression slope was examined in SPSS 23.0 (SPSS). The relative contribution of changes in soluble sugars or starch to NSCs responses was analysed using the RELAIMPO package in R.

Hedges' *d* is an estimate of the standardized mean difference not biased by small sample sizes (Gurevitch & Hedges, 2001). Thus, we employed Hedges' *d* to calculate interactive effects for each study according to the method of Crain, Kroeker, and Halpern (2008). For a study with two factors (A and B), main effect of factor A (d_A), main effect of factor B (d_B) and interactive effect (d_1) were calculated as following Equations 6–8:

$$d_{\rm A} = \frac{(X_{\rm AB} + X_{\rm A}) - (X_{\rm C} + X_{\rm B})}{2s} J(m), \tag{6}$$

$$d_{\rm B} = \frac{(X_{\rm AB} + X_{\rm B}) - (X_{\rm C} + X_{\rm A})}{2s} J(m), \tag{7}$$

$$d_{\rm I} = \frac{(X_{\rm AB} + X_{\rm C}) - (X_{\rm A} + X_{\rm B})}{2s} J(m), \tag{8}$$

where $X_{\rm C}$ is the mean value in the control group, and $X_{\rm A}$, $X_{\rm B}$ and $X_{\rm AB}$ are mean values in the experimental groups of A, B and their combination (A and B). J(m) and *s* represent correction term for small sample bias and pooled standard deviation respectively (Hedges & Olkin, 1985), which were estimated by the following Equations 9 and 10, respectively:

$$J(m) = 1 - \frac{3}{4m - 1},$$
(9)

$$s = \sqrt{\frac{(n_{\rm C} - 1)s_{\rm C}^2 + (n_{\rm A} - 1)s_{\rm A}^2 + (n_{\rm B} - 1)s_{\rm B}^2 + (n_{\rm AB} - 1)s_{\rm AB}^2}{n_{\rm C} + n_{\rm A} + n_{\rm B} + n_{\rm AB} - 4}},$$
 (10)

where *m* is the degree of freedom ($m = n_{\rm C} + n_{\rm A} + n_{\rm B} + n_{\rm AB} - 4$), $n_{\rm C}$, $n_{\rm A}$, $n_{\rm B}$ and $n_{\rm AB}$ are the sample sizes, and $s_{\rm C}$, $s_{\rm A}$, $s_{\rm B}$ and $s_{\rm AB}$ are the standard deviations in the corresponding groups. The variance of $d_1(v_2)$ was calculated by Equation 11:

$$v_{2} = \frac{1}{4} \left[\frac{1}{n_{C}} + \frac{1}{n_{A}} + \frac{1}{n_{B}} + \frac{1}{n_{AB}} + \frac{d_{I}^{2}}{2(n_{C} + n_{A} + n_{B} + n_{AB})} \right].$$
 (11)

The weight (w_2) can be calculated as reciprocal of variance (v_2). Likewise, it can be adjusted by the total number of observations per study (n) to calculate w'_2 (Li et al., 2018):

$$w_2 = \frac{1}{v_2},$$
 (12)

$$w_2' = \frac{w_2}{n}.$$
 (13)

The mean $d_1(d_{++})$ and standard error $[s(d_{++})]$ were estimated by Equations 14 and 15, respectively.

$$d_{++} = \frac{\sum_{i=1}^{l} \sum_{j=1}^{k} w'_{ij} d_{ij}}{\sum_{i=1}^{l} \sum_{k=1}^{k} w'_{ii}},$$
(14)

$$s(d_{++}) = \sqrt{\frac{1}{\sum_{i=1}^{l} \sum_{j=1}^{k} w_{ij}'}},$$
(15)

where *l* is the number of groups, *k* is the number of comparisons in the *i*th group. The 95% Cl of d_{++} was estimated as $d_{++} \pm C_{\partial/2} \times s(d_{++})$, where $C_{\partial/2}$ is the two-tailed critical value of the standard normal distribution. If the sample size was lower than 20, a bootstrapping method based on 9,999 iterations was used.

The interactive effect between two environmental change drivers was classified as additive, synergistic or antagonistic (Crain et al., 2008). If the 95% confidence interval (CI) overlapped with zero, then their interactive impact was additive. For the non-additive interactions (synergistic or antagonistic), there were three cases: (a) For two-driver pairs whose individual impacts were both negative, then their interactive impacts lower than zero was synergistic and greater than zero was antagonistic. (b) In cases where the individual effects were both positive, their interactive impact greater than zero was synergistic and lower than zero was antagonistic. (c) If the individual main impact was negative for one driver but positive for the other driver, the type of interactive effect was jointly determined by the arithmetic sum of the two individual main impacts and the calculated interactive effect by Equation 8 (Zhou et al., 2016). If the arithmetic sum of the two drivers' main effects (e.g. 0.36 for the $eCO_2 \times$ warming on sugars) and their interactive effect (e.g. 0.36 for the e $CO_2 \times$ warming on sugars) were both positive or negative, then their interactive effect was defined as synergistic. Otherwise, the interactive effect was defined as antagonistic.

3 | RESULTS

3.1 | Main effects of global change drivers on NSCs and its components

Elevated CO₂ concentration significantly increased NSCs (+27.8%), soluble sugars (+22.2%) and starch (+43.8%), and the enhancement of starch was nearly two times as large as that of soluble sugars. Nitrogen addition induced significant increases in NSCs (+4.1%), soluble sugars (+6.0%) and starch (+1.2%). NSCs (-4.1%) and starch (-8.5%) were significantly reduced under drought, yet soluble sugars were increased by 4.5%. Warming significantly reduced NSCs (-8.7%), soluble sugars (-7.0%) and starch (-10.2%; Figure 1a).

We further examined the relative impacts of soluble sugars and starch on the total NSCs in response to global-change drivers. The response ratio of NSCs exhibited positive correlations with response ratio of soluble sugars or starch under four different environmental changes (Figure 2). As shown by the difference in correlation slopes in Figure 2, the changes in response of NSCs to N addition and drought were more sensitive to that of soluble sugars than starch (both p < 0.01). Moreover, the relative contribution analysis showed that NSCs were mainly driven by soluble sugars under N addition (52%) and drought (76%), but by starch under eCO₂ (57%) and warming (92%; Figure 2).

Stronger enhancement of starch than soluble sugars under eCO_2 was found in woody and herbaceous plants as well as most plant functional types (Figure 3a; Table S1). Under N addition, starch

was increased more than soluble sugars in woody plants, whereas it was contrary to herbaceous plants. The different responses between soluble sugars and starch were found in broadleaved trees, forbs and annual herbs under N addition (Figure 3b; Table S1). The drought-induced different response between soluble sugars and starch was common except for conifers, deciduous trees and CAM herbs (Figure 3c; Table S1). Severe reduction of starch than soluble sugars under warming was found mainly in herbaceous species, such as forbs, perennial herbs, C_3 herbs (Figure 3d; Table S1).

Moreover, the responses of soluble sugars and starch under environmental changes varied with tissue types (Figure 4). Soluble sugars and starch were stimulated most in leaves under eCO₂. The significant difference between soluble sugars and starch was found in leaves and roots of woody plants as well as leaves and stems of herbaceous plants (Figure 4; Table S2). In woody plants, soluble sugars and starch were decreased in leaves (-9.2%) and roots (-19.9%) under N addition. In herbaceous plants, soluble sugars (+4.3%) were increased but starch (-28.4%) was decreased in leaves under N addition. Then the different response between starch and soluble sugars was only found in herbaceous leaves (Figure 4; Table S2). In leaves of woody plants and roots of herbaceous plants, soluble sugars were increased and starch was decreased under drought. Drought resulted in different responses between soluble sugars and starch in leaves and roots (Figure 4; Table S2). Decrease of soluble sugars was found in leaves and stems of woody plants and leaves and roots of herbaceous plants under warming. However, decrease of starch was only found in leaves of woody and herbaceous species. The different response between soluble sugars and starch was significant in the leaves of herbs (Figure 4; Table S2).

Soluble sugars (+27.2%) and starch (+53.8%) were enhanced most under moderate and low eCO_2 treatment respectively. The different response between soluble sugars and starch disappeared under high eCO_2 treatment (Figure 5a; Table S2). Soluble sugars (+19.7%) and starch (+2.1%) were increased under low N addition treatment. No significant difference between soluble sugars and starch was observed under N addition (Figure 5b; Table S2). Soluble sugars (+14.8%) were stimulated most under high drought treatment, yet starch (-24.3%) was decreased most under high drought treatment. The different response between them was found across the three intensities (Figure 5c; Table S2). Soluble sugars and starch were significantly decreased across the three intensities of warming,

FIGURE 1 Mean effect size of nonstructural carbohydrates (NSCs), soluble sugar (sugars) and starch under individual (a) and interactive (b) effects of multiple global change drivers. The error bars indicated the 95% confidence interval (Cl). If the Cl did not overlap with zero, a response was considered to be significant. The sample size for each variable is shown in the right column of the figure





FIGURE 2 Correlations of response ratio of NSCs with response ratios of its components (soluble sugars and starch) under (a) eCO_2 , (b) N addition, (c) drought and (d) warming. The pies represent the relative contributions of response ratios of soluble sugar and starch to responses ratio of NSCs. Points represent single pairs of data

FIGURE 3 Mean effect size of NSCs, soluble sugars and starch in different plant functional types under elevated CO₂ (a), N addition (b), drought (c) and warming (d). The error bars indicated 95% confidence interval (CI). If the CI did not overlap with zero, a response was considered to be significant. The sample size for each variable is shown in the same colour in the left and right column of the figure

FIGURE 4 Mean effect size of soluble sugars and starch in woody (a, c, e) and herbaceous (b, d, f) tissues (i.e. leaves, stems and roots). If the confidence interval did not overlap with zero, a response was considered to be significant. The sample size for each variable is shown near the dots



FIGURE 5 Mean effect size of soluble sugars and starch categorized by experimental intensities (i.e. low, moderate and high) under elevated CO_2 (a), N addition (b), drought (c) and warming (d). If the confidence interval did not overlap with zero, a response was considered to be significant. The sample size for each variable is shown near the dots

especially under high warming treatment. Low or high warming treatment induced a significant difference between soluble sugars and starch (Figure 5d; Table S2).

The increases of starch under eCO_2 and soluble sugars under drought decreased with the experimental duration (Figure S3). However, under N addition and warming, the effect size was independent of the experimental duration. The difference between soluble sugars and starch was found in both the field and laboratory conditions under eCO_2 and drought. However, it was significant only under field warming and insignificant under either natural or laboratory N addition experiment (Figure S5). Moreover, the response ratio of soluble sugars or starch was irrelative to mean annual temperature (Figure S6).

3.2 | Interactive effects of environmental change drivers on soluble sugars and starch

The interactive effects of eCO_2 with drought, N with drought, N with warming as well as drought with warming were all additive (Figure 1b). The mean effect size of soluble sugars (0.36) was greater than zero under eCO_2 with warming. The sum of main effects of eCO_2 and warming (0.56) on soluble sugars was larger than zero.

Hence, their interactive effect was synergistic (Figure 1b; Figure S2). The mean effect size of starch was lower than zero under eCO_2 with N addition (-0.44) or eCO_2 with warming (-0.40). However, the sum of main effects of eCO_2 and N addition (0.52) or eCO_2 and warming (0.78) on starch was larger than zero. Hence, their interactive effects were both antagonistic (Figure 1b; Figure S2).

The interactive effects of eCO_2 with drought, eCO_2 with warming, nitrogen with drought and nitrogen with warming were unrelated to experimental duration. However, the interactive effect of eCO_2 with N addition and drought with warming on starch showed a positive and negative relationship with experimental duration respectively (both p < 0.01; Figure S4).

4 | DISCUSSION

4.1 | Main effects of environmental change factors on NSCs and its components

Our study found that soluble sugars and starch reacted differently under four environmental changes. For example, starch was stimulated more than soluble sugars under eCO₂, especially in leaves (Figure 4). It is in accordance with results of Li et al. (2018), which has found eCO₂ increased starch and soluble sugars in leaves of trees by 125.3% and 20.5% respectively. Although leaf stomatal conductance is lowered under eCO_2 (Medlyn et al., 2011), the rate of carbon assimilation is still increased (Delucia, Sasek, & Strain, 1985; Long & Drake, 1992; Moore, Palmquist, & Seemann, 1997). Soluble sugars have higher metabolic activities and excessive sugars in the cytoplasm cause higher respiration rates and declined growth (Chapin III et al., 1990; Schulze, Stitt, Schulze, Neuhaus, & Fichtner, 1991). Moreover, cells with high sugar concentration are susceptible to osmotic dehydration (Ahmed, Qazi, & Jamal, 2016). In contrast, respiration is less affected by starch than sugars under eCO₂ because starch is insoluble and a long-term storage compound which is not readily available to participate in plant metabolic processes (Chapin III et al., 1990; Hoch, Richter, & Körner, 2003). We also found more enhancement of starch than soluble sugars across the three treatment intensities under CO₂ enhancement. However, starch was enhanced most under low or short-term CO₂ enhancement treatment (Figure 5a; Figure S3a). It suggests this stimulation is not persistent and limited by other factors, such as, N (Reich et al., 2014).

It has been suggested that the NSCs were significantly decreased in trees and herbs under N addition (Liu et al., 2016; Figure 4). Also, our meta-analysis found that starch decreased by 15.2% in the roots of trees under N addition. N addition commonly increases leaf photosynthesis and plant growth (Hyvonen et al., 2007; Lambers, Chapin III, & Pons, 2008; Nakaji, Fukami, Dokiya, & Izuta, 2001; Xia & Wan, 2008). For example, Xia and Wan (2008) have found that biomass and N concentration in plants are commonly increased by N addition. This increase in N concentration could be expected to be correlated with greater leaf respiration, based on the large-scale leaf-trait coordination. Hence, the consumption of stored starch could be enhanced to meet the increased carbohydrates demand for maintaining respiratory activities. Also, starch was significantly decreased under moderate or high N addition, which may arise from the elevated stimulation on root respiration and biomass along N addition gradients (Figure 4e; Li et al., 2015).

Our study detected that starch was significantly reduced, but soluble sugars were increased under drought (Figure 1a), consistent with the results of a recent data synthesis (Adams et al., 2017). It can occur by interconversion between soluble sugars and starch to meet osmoregulatory and metabolic demands of plants (McDowell, 2011). However, starch was not decreased and the NSCs were increased in herbs (Figure 3). Compared with woody species, herbaceous plants, especially herbaceous angiosperms, have similar abilities to avoid drought-induced embolism (Lens et al., 2016). Furthermore, herbaceous plants have more strategies to cope with water deficit (Volaire, 2018). For example, annual herbs with short growing seasons can end growth to escape water deficit (Kooyers, 2015). Herbs can coordinate plasticity of xylem to maintain hydraulic safety, such as, sunflowers (Cardoso, Brodribb, Lucani, DaMatta, & McAdam, 2018). Hence, herbaceous species are more tolerant to drought (Volaire, 2018). The discrepant utilization strategies of soluble sugars and starch were found in leaves and roots (Figure 4). These two organs are especially sensitive to water

availability and more starch will be converted into sugars in order to repair embolism and avoid catastrophic xylem failure (Sala, Woodruff, & Meinzer, 2012). Owing to stronger inhibition of photosynthesis under severe drought, the enhancement of soluble sugars disappeared, and starch was severely depleted in the long-term (Martínez-Vilalta et al., 2016; McDowell, 2011).

Warming-induced reductions of soluble sugars and starch in plants were common (Zvereva & Kozlov, 2006; Figures 1 and 3). It has been reported that rising temperature stimulates sucrose phosphate synthase (Dai et al., 2015; Hussain, Allen, & Bowes, 1999; Stitt & Grosse, 1988) and the starch-consuming α -amylase (Yamakawa, Hirose, Kuroda, & Yamaguchi, 2007). However, the activity of several enzymes involved in starch biosynthesis is reduced by warming (Hurkman et al., 2003; Jiang, Dian, & Wu, 2003; Wilhelm, Mullen, Keeling, & Singletary, 1999; Yamakawa et al., 2007). Furthermore, climate warming at night leads to a greater depletion of starch which can even in turn accelerate the accumulation of soluble sugars in the following day with enhanced leaf photosynthesis (Turnbull, Murthy, & Griffin, 2002; Turnbull et al., 2004; Wan, Xia, Liu, & Niu, 2009). Warming accelerates starch remobilization to provide energy for other activities and tissues (Thalmann & Santelia, 2017). Moreover, the positive response of respiration is enlarged with enhancing warming intensity (Liang, Xia, Liu, & Wan, 2013). Hence, starch was reduced most under high warming treatment (Figure 5). Effects of warming on plants are different among climate types, and different response between soluble sugars and starch was only found under field warming (Figure S5). It may because laboratory experiments have benign conditions for plant growth, such as temperature, water and nutrients (Poorter et al., 2016). However, response ratio of soluble sugars (p = 0.63) or starch (p = 0.92) was irrelative to mean annual temperature (Figure S6). The different functions performed by soluble sugars under drought (osmoregulation and metabolism mainly) and warming (metabolism mainly) explain discrepant relative contribution of response ratio of soluble sugars to response ratio of NSCs (Figure 2).

4.2 | Interactive effects of environmental change drivers on soluble sugars and starch

In our study, we found that additive effects on soluble sugars and starch were common between different global-change factors (Figure 1b). It is in accordance with the findings of Li et al. (2018). CO_2 enhancement and N addition significantly increased and decreased starch respectively (Figure S2). However, the interaction of eCO_2 with N addition reduced starch, suggesting that N is critical in limiting the positive impact of eCO_2 on starch accumulation (Wong, 1990). Considering starch is a long-term storage to support plant growth, such N limitation might be an important driver of the widely reported progressive N limitation on canopy growth and ecosystem productivity under eCO_2 (e.g. Luo et al., 2004; Norby, Warren, Iversen, Medlyn, & McMurtrie, 2010; Reich et al., 2014). The significant increase of soluble sugars and decrease of starch under eCO_2 with warming could result from more C allocation to immediate metabolism (mainly respiration and growth; Hussain et al., 1999; Liang et al., 2013) or decreased photosynthetic carbon accumulation under higher temperature (Aranjuelo, Irigoyen, Sánchez-Díaz, & Nogués, 2008). The antagonistic effects of eCO_2 with N addition and the additive effects of drought with warming showed significant relationships with experimental duration (Figure S4a,f). It implies these interactions may be changed in the long term.

5 | CONCLUSIONS

Our study quantifies the main and interactive effects of multiple environmental change factors on NSCs and its components. It reveals that (a) dynamics of NSCs are driven by starch under eCO₂ and warming, but by soluble sugars under N addition and drought; (b) Antagonistic interactions of eCO₂ with N addition and eCO2 with warming on starch were found. Synergistic interaction of eCO₂ with warming was found on soluble sugars; (c) Responses of soluble sugars and starch are closely related to plant functional types, tissue types and experimental conditions (i.e. treatment intensities, durations and approaches). Although additive interactions are dominant across two-pair drivers, it should be noted that experiments including interactions among three or more factors are still scarce. The discrepant responses of soluble sugars and starch under varying environmental conditions indicate that their different ecological functions need to be considered in canopy photosynthesis models (Dietze et al., 2014; Richardson et al., 2015). Thus, the consideration of NSCs' dynamics in vegetation models could be important under future global environmental changes, especially with the increasing frequency and intensity of extreme climate events such as heat waves and drought which will drastically alter the dynamics of NSCs in plants (Hansen, Sato, & Ruedy, 2012; Meehl & Tebaldi, 2004; Shi, Kloog, Zanobetti, Liu, & Schwartz, 2015).

ACKNOWLEDGEMENTS

We thank three anonymous reviewers for constructive comments to considerably improve the manuscript. We are grateful to all the authors who have contributed data to this analysis. This research was financially supported by National Natural Science Foundation of China (31722009, 41630528) and the National 1000 Young Talents Program of China.

AUTHORS' CONTRIBUTIONS

J.X. designed the study and Y.D. conducted the analysis; R.L. contributed to the data collection and discussion of results; J.X. and Y.D. wrote the manuscript. The authors declare no competing financial interests.

DATA AVAILABILITY STATEMENT

Data are deposited in the Dryad Digital Repository: https://doi. org/10.5061/dryad.j6q573n9n (Du, Lu, & Xia, 2020).

ORCID

Ying Du b https://orcid.org/0000-0001-9690-4757 Ruiling Lu https://orcid.org/0000-0001-9583-550X Jianyang Xia https://orcid.org/0000-0001-5923-6665

REFERENCES

- Adams, H. D., Zeppel, M. J. B., Anderegg, W. R. L., Hartmann, H., Landhäusser, S. M., Tissue, D. T., ... McDowell, N. G. (2017). A multi-species synthesis of physiological mechanisms in droughtinduced tree mortality. *Nature Ecology & Evolution*, 1, 1285–1291. https://doi.org/10.1038/s41559-017-0248-x
- Agami, R. A., Alamri, S. A., El-Mageed, T. A., Abousekken, M. S. M., & Hashem, M. (2018). Role of exogenous nitrogen supply in alleviating the deficit irrigation stress in wheat plants. *Agricultural Water Management*, 210, 261–270. https://doi.org/10.1016/j.agwat.2018.08.034
- Ahmed, I., Qazi, I. M., & Jamal, S. (2016). Developments in osmotic dehydration technique for the preservation of fruits and vegetables. *Innovative Food Science & Emerging Technologies*, 34, 29–43. https:// doi.org/10.1016/j.ifset.2016.01.003
- Aranjuelo, I., Irigoyen, J. J., Sánchez-Díaz, M., & Nogués, S. (2008). Carbon partitioning in N₂ fixing *Medicago sativa* plants exposed to different CO₂ and temperature conditions. *Functional Plant Biology*, 35, 306–317. https://doi.org/10.1071/FP07296
- Atkin, O. K., & Tjoelker, M. G. (2003). Thermal acclimation and the dynamic response of plant respiration to temperature. *Trends in Plant Science*, 8(7), 343–351. https://doi.org/10.1016/S1360-1385(03)00136-5
- Borenstein, M., Hedges, L. V., Higgins, J. P. T., & Rothstein, H. R. (2009). Introduction to meta-analysis. Chichester, UK: John Wiley & Sons Ltd. https://doi.org/10.1002/9780470743386
- Borenstein, M., Hedges, L. V., Higgins, J. P. T., & Rothstein, H. R. (2010). A basic introduction to fixed-effect and random-effects models for meta-analysis. *Research Synthesis Methods*, 1, 97–111. https://doi. org/10.1002/jrsm.12
- Brüggemann, N., Gessler, A., Kayler, Z., Keel, S. G., Badeck, F., Barthel, M., ... Bahn, M. (2011). Carbon allocation and carbon isotope fluxes in the plant-soil-atmosphere continuum: A review. *Biogeosciences*, *8*, 3619–3695. https://doi.org/10.5194/bgd-8-3619-2011
- Cardoso, A. A., Brodribb, T. J., Lucani, C. J., DaMatta, F. M., & McAdam, S. A. (2018). Coordinated plasticity maintains hydraulic safety in sunflower leaves. *Plant, Cell & Environment*, 41, 2567–2576. https://doi. org/10.1111/pce.13335
- Chapin III, F. S., Schulze, E. D., & Mooney, H. A. (1990). The ecology and economics of storage in plants. *Annual Review of Ecology and Systematics*, 21, 423–447. https://doi.org/10.1146/annurev.ecols ys.21.1.423
- Chaves, M. M., Maroco, J. P., & Pereira, J. S. (2003). Understanding plant responses to drought-from genes to the whole plant. *Functional Plant Biology*, 30, 239–264. https://doi.org/10.1071/FP02076
- Crain, C. M., Kroeker, K., & Halpern, B. S. (2008). Interactive and cumulative effects of multiple human stressors in marine systems. *Ecology Letters*, 11, 1304–1315. https://doi.org/10.1111/j.1461-0248.2008. 01253.x
- Dai, Y., Chen, B., Meng, Y., Zhao, W., Zhou, Z., Oosterhuis, D. M., & Wang, Y. (2015). Effects of elevated temperature on sucrose metabolism and cellulose synthesis in cotton fibre during secondary cell wall development. *Functional Plant Biology*, 42, 909–919. https://doi. org/10.1071/FP14361
- De Schepper, V., De Swaef, T., Bauweraerts, I., & Steppe, K. (2013). Phloem transport: A review of mechanisms and controls. *Journal of Experimental Botany*, 64, 4839–4850. https://doi.org/10.1093/jxb/ert302
- Delucia, E. H., Sasek, T. W., & Strain, B. R. (1985). Photosynthetic inhibition after long-term exposure to elevated levels of atmospheric

carbon dioxide. Photosynthesis Research, 7, 175–184. https://doi. org/10.1007/BF00037008

- Dietze, M. C., Sala, A., Carbone, M. S., Czimczik, C. I., Mantooth, J. A., Richardson, A. D., & Vargas, R. (2014). Nonstructural carbon in woody plants. Annual Review of Plant Biology, 65, 667–687. https:// doi.org/10.1146/annurev-arplant-050213-040054
- Du, Y., Lu, R., & Xia, J. (2020). Data from: Impacts of global environmental change drivers on non-structural carbohydrates in terrestrial plants. *Dryad Digital Repository*, https://doi.org/10.5061/ dryad.j6q573n9n
- Duan, H., Amthor, J. S., Duursma, R. A., O'Grady, A. P., Choat, B., & Tissue, D. T. (2013). Carbon dynamics of eucalypt seedlings exposed to progressive drought in elevated [CO₂] and elevated temperature. *Tree Physiology*, 33, 779–792. https://doi.org/10.1093/treephys/tpt061
- Gibon, Y., Pyl, E. T., Sulpice, R., Lunn, J. E., Hoehne, M., Guenther, M., & Stitt, M. (2009). Adjustment of growth, starch turnover, protein content and central metabolism to a decrease of the carbon supply when *Arabidopsis* is grown in very short photoperiods. *Plant, Cell & Environment*, 32, 859–874. https://doi.org/10.1111/j.1365-3040.2009.01965.x
- Gurevitch, J., & Hedges, L. V. (2001). Meta-analysis: Combining the results of independent experiments. In S. Scheiner & J. Gurevitch (Eds.), *Design and analysis of ecological experiments* (pp. 347–369). New York, NY: Oxford University Press.
- Habermann, E., San Martin, J. A. B., Contin, D. R., Bossan, V. P., Barboza, A., Braga, M. R., ... Martinez, C. A. (2019). Increasing atmospheric CO_2 and canopy temperature induces anatomical and physiological changes in leaves of the C_4 forage species *Panicum maximum*. *PLoS ONE*, 14, e0212506. https://doi.org/10.1371/journal.pone.0212506
- Hansen, J., Sato, M., & Ruedy, R. (2012). Perception of climate change. Proceedings of the National Academy of Sciences of the United States of America, 109, E2415–E2423. https://doi.org/10.1073/pnas.12052 76109
- Hartmann, H., & Trumbore, S. (2016). Understanding the roles of nonstructural carbohydrates in forest trees – From what we can measure to what we want to know. New Phytologist, 211, 386–403. https:// doi.org/10.1111/nph.13955
- Hedges, L. V., Gurevitch, J., & Curtis, P. S. (1999). The meta-analysis of response ratios in experimental ecology. *Ecology*, 80, 1150–1156. https://doi.org/10.2307/177062
- Hedges, L. V., & Olkin, I. (1985). *Statistical method for meta-analysis*. New York, NY: Academic Press.
- Hoch, G., Richter, A., & Körner, C. (2003). Non-structural carbon compounds in temperate forest trees. *Plant, Cell & Environment, 26*, 1067– 1081. https://doi.org/10.1046/j.0016-8025.2003.01032.x
- Huang, K., Xia, J., Wang, Y., Ahlström, A., Chen, J., Cook, R. B., ... Luo, Y. (2018). Enhanced peak growth of global vegetation and its key mechanisms. *Nature Ecology & Evolution*, *2*, 1897–1905. https://doi. org/10.1038/s41559-018-0714-0
- Hughes, L. (2000). Biological consequences of global warming: Is the signal already apparent? Trends in Ecology & Evolution, 15, 56–61. https:// doi.org/10.1016/s0169-5347(99)01764-4
- Hurkman, W. J., McCue, K. F., Altenbach, S. B., Korn, A., Tanaka, C. K., Kothari, K. M., ... DuPont, F. M. (2003). Effect of temperature on expression of genes encoding enzymes for starch biosynthesis in developing wheat endosperm. *Plant Science*, 164, 873–881. https://doi. org/10.1016/S0168-9452(03)00076-1
- Hussain, M. W., Allen Jr., L. H., & Bowes, G. (1999). Up-regulation of sucrose phosphate synthase in rice grown under elevated CO₂ and temperature. *Photosynthesis Research*, 60, 199–208. https://doi. org/10.1023/A:1006242001390
- Huttunen, L., Saravesi, K., Markkola, A., & Niemelä, P. (2013). Do elevations in temperature, CO₂, and nutrient availability modify belowground carbon gain and root morphology in artificially defoliated silver birch seedlings?*Ecology and Evolution*, *3*, 2783–2794. https:// doi.org/10.1002/ece3.665

- Hyvonen, R., Agren, G. I., Linder, S., Persson, T., Cotrufo, M. F., Ekblad, A., ... Wallin, G. (2007). The likely impact of elevated [CO₂], nitrogen deposition, increased temperature, and management on carbon sequestration in temperate and boreal forest ecosystems: A literature review. *New Phytologist*, 173(3), 463–480. https://doi. org/10.1111/j.1469-8137.2007.01967.x
- Jacobsen, K. R., Fisher, D. G., Maretzki, A., & Moore, P. H. (1992). Developmental changes in the anatomy of the sugarcane stem in relation to phloem unloading and sucrose storage. *Botanica Acta*, 105, 70–80. https://doi.org/10.1111/j.1438-8677.1992.tb00269.x
- Jiang, H., Dian, W., & Wu, P. (2003). Effect of high temperature on fine structure of amylopectin in rice endosperm by reducing the activity of the starch branching enzyme. *Phytochemistry*, 63, 53–59. https:// doi.org/10.1016/S0031-9422(03)00005-0
- Kooyers, N. J. (2015). The evolution of drought escape and avoidance in natural herbaceous populations. *Plant Science*, 234, 155–162. https:// doi.org/10.1016/j.plantsci.2015.02.012
- Körner, C. (2003). Carbon limitation in trees. *Journal of Ecology*, 91, 4–17. https://doi.org/10.1046/j.1365-2745.2003.00742.x
- Kozlowski, T. T. (1992). Carbohydrate sources and sinks in woody plants. Botanical Review, 58(2), 107–222. https://doi.org/10.1007/BF028 58600
- Lambers, H., Chapin III, F. S., & Pons, T. L. (2008). Plant physiological ecology (2nd ed.). New York, NY: Springer-Verlag. https://doi. org/10.1093/aob/mcn225
- Landhäusser, S. M., Chow, P. S., Dickman, L. T., Furze, M. E., Kuhlman, I., Schmid, S., ... Adams, H. D. (2018). Standardized protocols and procedures can precisely and accurately quantify non-structural carbohydrates. *Tree Physiology*, 38(12), 1764–1778. https://doi.org/10.1093/ treephys/tpy118
- Lens, F., Picon-Cochard, C., Delmas, C. E. L., Signarbieux, C., Buttler, A., Cochard, H., ... Delzon, S. (2016). Herbaceous angiosperms are not more vulnerable to drought-induced embolism than angiosperm trees. *Plant Physiology*, 172, 661–667. https://doi.org/10.1104/pp.16. 00829
- Li, W., Hartmann, H., Adams, H. D., Zhang, H., Jin, C., Zhao, C., ... Wu, J. (2018). The sweet side of global change-dynamic responses of non-structural carbohydrates to drought, elevated CO₂ and nitrogen fertilization in tree species. *Tree Physiology*, 38, 1706–1723. https:// doi.org/10.1093/treephys/tpy059
- Li, W., Jin, C., Guan, D., Wang, Q., Wang, A., Yuan, F., & Wu, J. (2015). The effects of simulated nitrogen deposition on plant root traits: A meta-analysis. *Soil Biology and Biochemistry*, 82, 112–118. https://doi. org/10.1016/j.soilbio.2015.01.001
- Liang, J., Xia, J., Liu, L., & Wan, S. (2013). Global patterns of the responses of leaf-level photosynthesis and respiration in terrestrial plants to experimental warming. *Journal of Plant Ecology*, *6*, 437–447. https://doi. org/10.1093/jpe/rtt003
- Liu, J., Wu, N., Wang, H., Sun, J., Peng, B., Jiang, P., & Bai, E. (2016). Nitrogen addition affects chemical compositions of plant tissues, litter and soil organic matter. *Ecology*, 97, 1796–1806. https://doi. org/10.1890/15-1683.1
- Long, S. P., & Drake, B. G. (1992). Photosynthetic CO₂ assimilation and rising atmospheric CO₂ concentrations. In N. Baker & H. Thomas (Eds.), Crop photosynthesis: Spatial and temporal determinants (pp. 69–103). Amsterdam, The Netherlands: Elsevier Science Publishers.
- Luo, Y., Su, B. O., Currie, W. S., Dukes, J. S., Finzi, A., Hartwig, U., ... Pataki, D. E. (2004). Progressive nitrogen limitation of ecosystem responses to rising atmospheric carbon dioxide. *BioScience*, 54, 731–739. https://doi.org/10.1641/0006-3568(2004)054[0731:pnloer]2.0. co;2
- Martínez-Vilalta, J., Sala, A., Asensio, D., Galiano, L., Hoch, G., Palacio, S., ... Lloret, F. (2016). Dynamics of non-structural carbohydrates in terrestrial plants: A global synthesis. *Ecological Monographs*, 86, 495– 516. https://doi.org/10.1002/ecm.1231

- McDowell, N. G. (2011). Mechanisms linking drought, hydraulics, carbon metabolism, and vegetation mortality. *Plant Physiology*, 155, 1051– 1059. https://doi.org/10.1104/pp.110.170704
- McDowell, N. G., & Allen, C. D. (2015). Darcy's law predicts widespread forest mortality under climate warming. *Nature Climate Change*, 5, 669–672. https://doi.org/10.1038/nclimate2641
- Medlyn, B. E., Duursma, R. A., Eamus, D., Ellsworth, D. S., Prentice, I. C., Barton, C. V. M., ... Wingate, L. (2011). Reconciling the optimal and empirical approaches to modelling stomatal conductance. *Global Change Biology*, *17*, 2134–2144. https://doi.org/10.1111/j.1365-2486. 2010.02375.x
- Meehl, G. A., & Tebaldi, C. (2004). More intense, more frequent, and longer lasting heat waves in the 21st century. *Science*, 305, 994–997. https://doi.org/10.1126/science.1098704
- Moore, B. D., Palmquist, D. E., & Seemann, J. R. (1997). Influence of plant growth at high CO₂ concentrations on leaf content of ribulose-1,5-bisphosphate carboxylase/oxygenase and intracellular distribution of soluble carbohydrates in tobacco, snapdragon, and parsley. *Plant Physiology*, 115, 241–248. https://doi.org/10.1104/ pp.115.1.241
- Nakaji, T., Fukami, M., Dokiya, Y., & Izuta, T. (2001). Effects of high nitrogen load on growth, photosynthesis and nutrient status of *Cryptomeria japonica* and *Pinus densiflora* seedlings. *Trees*, 15, 453– 461. https://doi.org/10.1007/s00468-001-0130-x
- Norby, R. J., Warren, J. M., Iversen, C. M., Medlyn, B. E., & McMurtrie, R. E. (2010). CO₂ enhancement of forest productivity constrained by limited nitrogen availability. *Proceedings of the National Academy of Sciences of the United States of America*, 107, 19368–19373. https:// doi.org/10.1073/pnas.1006463107
- Parmesan, C., & Yohe, G. (2003). A globally coherent fingerprint of climate change impacts across natural systems. *Nature*, 421, 37-42. https://doi.org/10.1038/nature01286
- Plavcová, L., & Jansen, S. (2015). The role of xylem parenchyma in the storage and utilization of nonstructural carbohydrates. In U. Hacke (Ed.), *Functional and ecological xylem anatomy* (pp. 209–234). Cham, Switzerland: Springer.
- Poorter, H., Fiorani, F., Pieruschka, R., Wojciechowski, T., van derPutten, W. H., Kleyer, M., ... Postma, J. (2016). Pampered inside, pestered outside? Differences and similarities between plants growing in controlled conditions and in the field. *New Phytologist*, 212, 838–855. https://doi.org/10.1111/nph.14243
- Quentin, A. G., Pinkard, E. A., Ryan, M. G., Tissue, D. T., Baggett, L. S., Adams, H. D., ... Woodruff, D. R. (2015). Non-structural carbohydrates in woody plants compared among laboratories. *Tree Physiology*, 35, 1146–1165. https://doi.org/10.1093/treephys/tpv073
- R Development Core Team. (2016). R: A language and environment for statistical computing. Vienna, Austria: R Foundation for Statistical Computing. Retrieved from http://www.R-project.org/
- Rae, A. L., Perroux, J. M., & Grof, C. P. (2005). Sucrose partitioning between vascular bundles and storage parenchyma in the sugarcane stem: A potential role for the ShSUT1 sucrose transporter. *Planta*, 220, 817–825. https://doi.org/10.1007/s00425-004-1399-y
- Reich, P. B., Hobbie, S. E., & Lee, T. D. (2014). Plant growth enhancement by elevated CO₂ eliminated by joint water and nitrogen limitation. *Nature Geoscience*, 7, 920–924. https://doi.org/10.1038/ngeo2284
- Richardson, A. D., Carbone, M. S., Huggett, B. A., Furze, M. E., Czimczik, C. I., Walker, J. C., ... Murakami, P. (2015). Distribution and mixing of old and new nonstructural carbon in two temperate trees. *New Phytologist*, 206, 590–597. https://doi.org/10.1111/nph.13273
- Rolland, F., Baena-Gonzalez, E., & Sheen, J. (2006). Sugar sensing and signaling in plants: Conserved and novel mechanisms. *Annual Review of Plant Biology*, *57*, 675–709. https://doi.org/10.1146/annurev.arplant. 57.032905.105441
- Rosenberg, M. S., Adams, D. C., & Gurevitch, J. (2000). MetaWin: Statistical software for meta-analysis. Sunderland, MA: Sinauer Associates.

- Sala, A., Woodruff, D. R., & Meinzer, F. C. (2012). Carbon dynamics in trees: Feast or famine?Tree Physiology, 32, 764–775. https://doi. org/10.1093/treephys/tpr143
- Schulze, W., Stitt, M., Schulze, E. D., Neuhaus, H. E., & Fichtner, K. (1991). A quantification of the significance of assimilatory starch for growth of Arabidopsis thaliana L. Heynh. Plant Physiology, 95, 890–895. https://doi.org/10.1104/pp.95.3.890
- Shi, L., Kloog, I., Zanobetti, A., Liu, P., & Schwartz, J. D. (2015). Impacts of temperature and its variability on mortality in New England. *Nature Climate Change*, 5, 988–991. https://doi.org/10.1038/nclimate 2704
- Slewinski, T. L. (2012). Non-structural carbohydrate partitioning in grass stems: A target to increase yield stability, stress tolerance, and biofuel production. *Journal of Experimental Botany*, 63, 4647–4670. https://doi.org/10.1093/jxb/ers124
- Smith, S. E., & Smith, F. A. (2011). Roles of arbuscular mycorrhizas in plant nutrition and growth: New paradigms from cellular to ecosystem scales. Annual Review of Plant Biology, 62, 227–250. https://doi. org/10.1146/annurev-arplant-042110-103846
- Steele, C. L., Lewinsohn, E., & Croteau, R. (1995). Induced oleoresin biosynthesis in grand fir as a defense against bark beetles. Proceedings of the National Academy of Sciences of the United States of America, 92, 4164–4168. https://doi.org/10.1073/pnas.92.10.4164
- Stitt, M., & Grosse, H. (1988). Interactions between sucrose synthesis and CO₂ fixation IV. Temperature-dependent adjustment of the relation between sucrose synthesis and CO₂ fixation. *Journal of Plant Physiology*, 133, 392–400. https://doi.org/10.1016/S0176-1617(88)80025-7
- Sulpice, R., Flis, A., Ivakov, A. A., Apelt, F., Krohn, N., Encke, B., ... Stitt, M. (2014). Arabidopsis coordinates the diurnal regulation of carbon allocation and growth across a wide range of photoperiods. Molecular Plant, 7, 137–155. https://doi.org/10.1093/mp/sst127
- Thalmann, M., & Santelia, D. (2017). Starch as a determinant of plant fitness under abiotic stress. New Phytologist, 214, 943–951. https://doi. org/10.1111/nph.14491
- Thompson, S. G., & Higgins, J. P. (2002). How should meta-regression analyses be undertaken and interpreted?Statistics in Medicine, 21, 1559–1573. https://doi.org/10.1002/sim.1187
- Turnbull, M. H., Murthy, R., & Griffin, K. L. (2002). The relative impacts of daytime and night-time warming on photosynthetic capacity in *Populus deltoides. Plant, Cell & Environment, 25, 1729–1737. https://* doi.org/10.1046/j.1365-3040.2002.00947.x
- Turnbull, M. H., Tissue, D. T., Murthy, R., Wang, X., Sparrow, A. D., & Griffin, K. L. (2004). Nocturnal warming increases photosynthesis at elevated CO₂ partial pressure in *Populus deltoides*. New Phytologist, 161, 819–826. https://doi.org/10.1111/j.1469-8137.2004.00994.x
- Volaire, F. (2018). A unified framework of plant adaptive strategies to drought: Crossing scales and disciplines. *Global Change Biology*, 24, 2929–2938. https://doi.org/10.1111/gcb.14062
- Wan, S., Xia, J., Liu, W., & Niu, S. (2009). Photosynthetic overcompensation under nocturnal warming enhances grassland carbon sequestration. *Ecology*, 90, 2700–2710. https://doi.org/10.1890/08-2026.1
- Wilhelm, E. P., Mullen, R. E., Keeling, P. L., & Singletary, G. W. (1999). Heat stress during grain filling in maize: Effects on kernel growth and metabolism. *Crop Science*, *39*, 1733–1741. https://doi.org/10.2135/ cropsci1999.3961733x
- Wong, S. C. (1990). Elevated atmospheric partial pressure of CO₂ and plant growth. *Photosynthesis Research*, *23*(2), 171–180. https://doi. org/10.1007/BF00035008
- Xia, J., & Wan, S. (2008). Global response patterns of terrestrial plant species to nitrogen addition. New Phytologist, 179, 428–439. https:// doi.org/10.1111/j.1469-8137.2008.02488.x
- Xu, X., Yan, L., & Xia, J. (2019). A threefold difference in plant growth response to nitrogen addition between the laboratory and field experiments. *Ecosphere*, 10, e02572. https://doi.org/10.1002/ecs2.2572

- Xu, Z., Shimizu, H., Yagasaki, Y., Ito, S., Zheng, Y., & Zhou, G. (2013). Interactive effects of elevated CO₂, drought, and warming on plants. *Journal of Plant Growth Regulation*, 32, 692–707. https://doi. org/10.1007/s00344-013-9337-5
- Yamakawa, H., Hirose, T., Kuroda, M., & Yamaguchi, T. (2007). Comprehensive expression profiling of rice grain filling-related genes under high temperature using DNA microarray. *Plant Physiology*, 144, 258–277. https://doi.org/10.1104/pp.107.098665
- Yuan, Z. Y., & Chen, H. Y. (2015). Decoupling of nitrogen and phosphorus in terrestrial plants associated with global changes. *Nature Climate Change*, 5, 465–469. https://doi.org/10.1038/nclimate2549
- Yue, K., Fornara, D. A., Yang, W., Peng, Y., Li, Z., Wu, F., & Peng, C. (2017). Effects of three global change drivers on terrestrial C: N: P stoichiometry: A global synthesis. *Global Change Biology*, 23, 2450–2463. https://doi.org/10.1111/gcb.13569
- Yue, K., Fornara, D. A., Yang, W., Peng, Y., Peng, C., Liu, Z., & Wu, F. (2017). Influence of multiple global change drivers on terrestrial carbon storage: Additive effects are common. *Ecology Letters*, 20, 663–672. https://doi.org/10.1111/ele.12767
- Yue, K., Yang, W., Peng, Y., Peng, C., Tan, B. O., Xu, Z., ... Wu, F. (2018). Individual and combined effects of multiple global change drivers on terrestrial phosphorus pools: A meta-analysis. *Science of the Total Environment*, 630, 181–188. https://doi.org/10.1016/j.scito tenv.2018.02.213

- Zhou, L., Zhou, X., Shao, J., Nie, Y., He, Y., Jiang, L., ... Hosseini Bai, S. (2016). Interactive effects of global change factors on soil respiration and its components: A meta-analysis. *Global Change Biology*, 22, 3157–3169. https://doi.org/10.1111/gcb.13253
- Zhu, Z., Piao, S., Myneni, R. B., Huang, M., Zeng, Z., Canadell, J. G., ... Zeng, N. (2016). Greening of the Earth and its drivers. *Nature Climate Change*, 6, 791–795. https://doi.org/10.1038/nclimate3004
- Zvereva, E. L., & Kozlov, M. V. (2006). Consequences of simultaneous elevation of carbon dioxide and temperature for plant-herbivore interactions: A meta-analysis. *Global Change Biology*, 12, 27–41. https:// doi.org/10.1111/j.1365-2486.2005.01086.x

SUPPORTING INFORMATION

Additional supporting information may be found online in the Supporting Information section.

How to cite this article: Du Y, Lu R, Xia J. Impacts of global environmental change drivers on non-structural carbohydrates in terrestrial plants. *Funct Ecol.* 2020;34:1525– 1536. https://doi.org/10.1111/1365-2435.13577