Modern Physics Letters B Vol. 33, No. 31 (2019) 1950390 (19 pages) © World Scientific Publishing Company DOI: 10.1142/S0217984919503901

Multi-dark soliton solutions for the (2 + 1)-dimensional multi-component Maccari system

Tao Xu^{*,†}, Yong Chen^{*,‡,§} and Zhijun Qiao^{†,¶}

* School of Mathematical Sciences, Shanghai Key Laboratory of PMMP, East China Normal University, Shanghai 200062, China [†]School of Mathematical and Statistical Science, The University of Texas Rio Grande Valley, Edinburg, TX 78539, USA [‡]Department of Physics, Zhejiang Normal University, Jinhua 321004, China [§]ychen@sei.ecnu.edu.cn ¶zhijun.giao@utrqv.edu

> Received 13 January 2019 Accepted 9 July 2019 Published 8 November 2019

Based on reduction of the KP hierarchy, the general multi-dark soliton solutions in Gram type determinant forms for the (2+1)-dimensional multi-component Maccari system are constructed. Especially, the two component coupled Maccari system comprising of two component short waves and single-component long waves are discussed in detail. Besides, the dynamics of one and two dark-dark solitons are analyzed. It is shown that the collisions of two dark-dark solitons are elastic by asymptotic analysis. Additionally, the two dark-dark solitons bound states are studied through two different cases (stationary and moving cases). The bound states can exist up to arbitrary order in the stationary case, however, only two-soliton bound state exists in the moving case. Besides, the oblique stationary bound state can be generated for all possible combinations of nonlinearity coefficients consisting of positive, negative and mixed cases. Nevertheless, the parallel stationary and the moving bound states are only possible when nonlinearity coefficients take opposite signs.

Keywords: (2 + 1)-dimensional multi-component Maccari system; reduction of the KP hierarchy; multi-dark soliton solutions; bound states.

1. Introduction

The study for multi-component nonlinear systems has grown popular rapidly in recent years, and a lot of novel and interesting physical phenomenons can be generated by the nonlinear interaction of multiple waves.^{1–5} Additionally, a variety of complex systems usually involve more than one component, such as nonlinear

[§]Corresponding author.



T. Xu, Y. Chen & Z.-J. Qiao

optical fibers and Bose–Einstein condensates, etc. It is greatly necessary to extend the corresponding researches to multi-component systems.^{6–8} There always exist coupled effects that cross-phase modulation in the multi-component ones, and the corresponding various solutions cannot be correlated by Galileo transformation.⁹ Compared to the single-component systems,^{10,11} affluent exact solutions may appear in multi-component systems. In Ref. 12, some anomalous Peregrine solitons were constructed for the coupled Fokas–Lenells equations. Nth-order vector rational and semi-rational rogue waves for the three-component nonlinear Schrödinger (NLS) equations were given by Darboux transformation.¹³ Utilizing reduction of the KP hierarchy, hybrid solutions were exhibited in the Mel'nikov system.¹⁴ Furthermore, the interactional solutions including higher-order RWs interacting with multi-soliton (or multi-breather) were constructed in different multi-component systems.^{15,16}

As one of the hot topics in nonlinear science, dark solitons have been reported in many documents.^{17–20} There have been many different methods for constructing the dark soliton, such as the binary Darboux transformation,^{17,21,22} the algebraic– geometry reduction method,¹⁹ the dressing-Hirota method²⁰ and reduction of the KP hierarchy.^{18,23,24} The main goal of this work is to construct multi-dark soliton solutions for the multi-component Maccari system through reduction of the KP hierarchy. In 1983, the Kyoto school first developed the KP hierarchy reduction technique,²⁵ which has been widely applied to generate dark solitons,^{18,23,24} bright-dark mixed solitons,^{26–28} general rogue wave solutions^{29–31} and interactional solutions^{14,32} for a lot of integrable nonlinear systems.

In this paper, we focus on the multi-dark soliton solutions for the following (2 + 1)-dimensional N-component Maccari system, which includes N short wave components and one long wave component

 $v_u = \langle \Lambda \phi, \phi^* \rangle_x$,

$$i\phi_t + \phi_{xx} + v\phi = \mathbf{0}\,,\tag{1}$$

(2)

where

$$\Lambda = \operatorname{diag}(\sigma_1, \sigma_2, \dots, \sigma_{N-1}, \sigma_N),$$

$$\phi = (\phi^{(1)}, \phi^{(2)}, \dots, \phi^{(N-1)}, \phi^{(N)})^T,$$

$$\phi^* = (\phi^{(1)*}, \phi^{(2)*}, \dots, \phi^{(N-1)*}, \phi^{(N)*})^T,$$

 ϕ , **0** and ϕ^* denote the *N*-dimensional column vectors. $\sigma_l = \pm 1$, $\phi^{(l)} \equiv \phi^{(l)}(x, y, t)$ (l = 1, 2, ..., N) denote *N* different complex short wave amplitudes and $v \equiv v(x, y, t)$ is the real long wave amplitude. Besides, the symbol * denotes complex conjugation and the subscripted variables x, y and t are the corresponding partial differentiations. The symbol "diag" denotes the *N*-order diagonal matrix and \langle , \rangle stands for the inner product.

Choosing N = 2 in the above systems (1) and (2), the following two component coupled Maccari system including two short wave components and one long wave

$$i\phi_t^{(1)} + \phi_{xx}^{(1)} + v\phi^{(1)} = 0, \qquad (3)$$

$$i\phi_t^{(2)} + \phi_{xx}^{(2)} + v\phi^{(2)} = 0, \qquad (4)$$

$$v_y = (\sigma_1 \phi^{(1)} \phi^{(1)*} + \sigma_2 \phi^{(2)} \phi^{(2)*})_x, \qquad (5)$$

where $\sigma_k = \pm 1(k = 1, 2)$, $\phi^{(1)} \equiv \phi^{(1)}(x, y, t)$ and $\phi^{(2)} \equiv \phi^{(2)}(x, y, t)$ are the complex short wave amplitudes and $v \equiv v(x, y, t)$ is the real long wave amplitude.

Through a special reduction method, the coupled systems (3)–(5) were first derived by Maccari.³³ Under some special transformations, the above-mentioned two component coupled Maccari system can be reduced to our known nonlinear models. When y = x, the coupled systems (3)–(5) becomes two component coupled NLS equations.³⁴ Choosing $\phi^{(1)} = \phi^{(2)*}$, the Maccari system is reduced to a (2+1)-dimensional extension of the NLS equation.³⁵ If y = t, we can get the coupled long wave resonance equations.³⁶ The Maccari system can describe some motions of isolated waves which are localized in small parts of space in many fields, such as nonlinear optics, hydrodynamics, plasma physics.³⁷

There have been many papers reported on the coupled Maccari system. The Painlevé integrability and some special types of the localized excitations were constructed in Ref. 38. Some types of doubly periodic propagating wave patterns were generated by the variable separation approach.³⁹ In Ref. 40, dromion solutions for the Maccari system were generated. Different types of rational solutions including multi-rogue waves, lump solitons and interactions between the two were all given in the Maccari system by bilinear method.³⁷ Additionally, the bright solitons⁴¹ and bright-dark mixed solitons^{28,42} for the Maccari system were all studied by different methods. To the best of our knowledge, multi-dark soliton solutions for the coupled Maccari system (3)–(5) and the corresponding multi-component generalization (1) and (2) have never been reported up to now. Here, we will construct the multi-dark soliton solutions for the Maccari system by reduction of the KP hierarchy.

In this paper, the N-dark soliton solutions in Gram type determinant for the multi-component Maccari system are generated through reduction of the KP hierarchy. Furthermore, the two component coupled case (3)–(5) are discussed in detail for an example. Some dynamics and figures of one- and two dark-dark solitons are given. Utilizing the asymptotic analysis technique, the collisions between the two solitons are proved to be elastic. It can be shown that the two dark-dark solitons bound states possess two cases consisting of stationary and moving bound states. The bound states can exist up to arbitrary order in the stationary case, however, only two-soliton bound states exist in the moving case. Moreover, the oblique stationary bound state can be generated for all possible combinations of nonlinearity coefficients consist of positive, negative and mixed cases. Nevertheless, the parallel stationary and the moving bound states are only possible when nonlinearity coefficients take opposite signs.

The paper is organized as follows. In Sec. 2, the formulae of N-dark-dark soliton solutions for the two component coupled Maccari system (3)–(5) are given and proved. In Sec. 3, some dynamics of one- and two dark-dark solitons are discussed in detail. In Sec. 4, the dark-dark soliton bound states including stationary and moving cases are exhibited. In Sec. 5, the uniform formulae of N-dark solitons for the multi-component generalization (1) and (2) are constructed. The last section includes some conclusions and discussions.

N-Dark-Dark Soliton Solutions for the (2 + 1)-Dimensional Two Component Coupled Maccari System

Through the dependent variable transformation

[

$$\phi^{(1)} = k_1 e^{i\theta_1} \frac{h^{(1)}}{f}, \quad \phi^{(2)} = k_2 e^{i\theta_2} \frac{h^{(2)}}{f}, \quad v = 2(\ln f)_{xx}, \tag{6}$$

the two component coupled Maccari system (3)–(5) can be transformed to the following bilinear forms:

$$D_x^2 + i(D_t + 2\alpha_1 D_x)]h^{(1)} \cdot f = 0, \qquad (7)$$

$$[D_x^2 + i(D_t + 2\alpha_2 D_x)]h^{(2)} \cdot f = 0, \qquad (8)$$

$$D_x D_y f \cdot f = \sigma_1 k_1^2 (h^{(1)} h^{(1)*} - f^2) + \sigma_2 k_2^2 (h^{(2)} h^{(2)*} - f^2), \qquad (9)$$

where k_1 and k_2 are arbitrary real constants, f, $h^{(1)}$ and $h^{(2)}$ are all the functions of x, y and t, besides, f is a real function, $h^{(1)}$ and $h^{(2)}$ are two complex functions. Meanwhile, $\theta_i = \alpha_i x - \alpha_i^2 t + \beta_i(y)(i = 1, 2)$, where α_i are arbitrary real constants ($\alpha_1 \neq \alpha_2$) and $\beta_i(y)$ are arbitrary real functions of y. The operator D denotes the Hirota's bilinear differential operator, which can be defined as follows:

$$D_x^l D_y^m D_t^n f(x, y, t) \cdot g(x, y, t) = \left(\frac{\partial}{\partial x} - \frac{\partial}{\partial x'}\right)^l \left(\frac{\partial}{\partial y} - \frac{\partial}{\partial y'}\right)^m \left(\frac{\partial}{\partial t} - \frac{\partial}{\partial t'}\right)^n \\ \times f(x, y, t), \quad g(x', y', t')|_{x=x', y=y', t=t'}.$$
(10)

From the KP theory, the nonlinear models under investigation can be seen as some special reductions of the integrable equations in the KP hierarchy.^{7,18} Then the solutions that are bright, dark and rational solutions for some relative nonlinear systems can be derived from the τ function in the KP hierarchy under the appropriate reductions. Here, the multi-dark soliton solutions for the Maccari system (3)–(5) can be generated from the reduction of single KP hierarchy consisting of two copies of shifted singular points. **Lemma 2.1.** Considering the following bilinear equations in the KP hierar $chy^{18,23,24}$

$$(D_{x_1}^2 + 2aD_{x_1} - D_{x_2})\tau(k+1,l) \cdot \tau(k,l) = 0, \qquad (11)$$

$$\left(\frac{1}{2}D_{x_1}D_{x_{-1}} - 1\right)\tau(k,l)\cdot\tau(k,l) = -\tau(k+1,l)\tau(k-1,l)\,,\tag{12}$$

$$(D_{x_1}^2 + 2bD_{x_1} - D_{x_2})\tau(k, l+1) \cdot \tau(k, l) = 0, \qquad (13)$$

$$\left(\frac{1}{2}D_{x_1}D_{y_{-1}} - 1\right)\tau(k,l)\cdot\tau(k,l) = -\tau(k,l+1)\tau(k,l-1),$$
(14)

where k and l are arbitrary integers, a and b are two complex constants. The above bilinear equations (11)–(14) have the Gram determinant solutions

$$\tau(k,l) = |m_{(i,j)}(k,l)|_{1 \le i,j \le N}, \qquad (15)$$

here, the entries of the determinant $\tau(k,l)$ are given as

$$m_{i,j}(k,l) = c_{ij} + \int \varphi_i(k,l)\psi_j(k,l)dx_1, \qquad (16)$$

$$\varphi_i(k,l) = (p_i - a)^k (p_i - b)^l e^{\eta_i}, \qquad (17)$$

$$\psi_j(k,l) = \left(-\frac{1}{q_j+a}\right)^k \left(-\frac{1}{q_j+b}\right)^l e^{\zeta_j} , \qquad (18)$$

with

$$\eta_i = \frac{1}{p_i - a} x_{-1} + \frac{1}{p_i - b} y_{-1} + p_i x_1 + p_i^2 x_2 + \eta_{i0} ,$$

$$\zeta_j = \frac{1}{q_j + a} x_{-1} + \frac{1}{q_j + b} y_{-1} + q_j x_1 - q_j^2 x_2 + \zeta_{j0} ,$$

where c_{ij} , p_i , q_j , η_{i0} and ζ_{j0} are complex constants.

We should point out that, Eqs. (11) and (13) are the lowest-degree bilinear equation in the first modified KP hierarchy,²⁵ Eqs. (12) and (14) are the bilinear equations for the two dimensional Toda lattice.^{25,43} The validity of Lemma 1 has been proved in Eq. (6) in Ref. 18 and we omit the processes of proof here.

In order to get the N-dark-dark soliton solutions of the Maccari system (3)–(5), we should consider the reduction of the bilinear equations in Lemma 1. Choosing that a, b and x_2 are pure imaginary, x_1, x_{-1}, y_{-1} are real, and fixing the following relations $a = i\alpha_1, b = i\alpha_2, q_j = p_j^*, \zeta_{j0} = \eta_{j0}^*, c_{ji} = c_{ij} = \delta_{ij}$ (δ_{ij} is the Kronecker symbol), the following equalities can be given

$$\zeta_j = \eta_j^*, \quad \tau(k,l) = \tau^*(-k,-l).$$
 (19)

Through defining $f, h^{(1)}, h^{(2)}$ as

$$f = \tau(0,0), \quad h^{(1)} = \tau(1,0), \quad h^{(2)} = \tau(0,1),$$

$$h^{(1)*} = \tau(-1,0), \quad h^{(2)*} = \tau(0,-1),$$

(20)

the bilinear equations (11)–(14) can be rewritten as

$$(D_{x_1}^2 + 2i\alpha_1 D_{x_1} - D_{x_2})h^{(1)} \cdot f = 0, \qquad (21)$$

$$\left(\frac{1}{2}D_{x_1}D_{x_{-1}} - 1\right)f \cdot f = -h^{(1)}h^{(1)*}, \qquad (22)$$

$$(D_{x_1}^2 + 2i\alpha_2 D_{x_1} - D_{x_2})h^{(2)} \cdot f = 0, \qquad (23)$$

$$\left(\frac{1}{2}D_{x_1}D_{y_{-1}} - 1\right)f \cdot f = -h^{(2)}h^{(2)*}.$$
(24)

If we introduce these independent variable transformations

$$x_1 = x, \quad x_2 = it, \quad x_{-1} = -\frac{1}{2}\sigma_1 k_1^2 y, \quad y_{-1} = -\frac{1}{2}\sigma_2 k_2^2 y,$$
 (25)

namely

$$\partial_x = \partial_{x_1}, \quad \partial_t = i\partial_{x_2}, \quad \partial_y = -\frac{1}{2}\sigma_1 k_1^2 \partial_{x_{-1}} - \frac{1}{2}\sigma_2 k_2^2 \partial_{y_{-1}}, \tag{26}$$

the bilinear equations (21)–(24) can be directly calculated to recast to Eqs. (7)–(9). Considering both the τ functions in Eq. (20) and the variable transformations (25), one can directly get the *N*-dark-dark soliton solution in Gram type determinant forms.

Theorem 2.1. The N-dark-dark soliton solutions of the two component coupled Maccari system (3)–(5) can be written as

$$\phi^{(1)} = k_1 e^{i\theta_1} \frac{h^{(1)}}{f}, \quad \phi^{(2)} = k_2 e^{i\theta_2} \frac{h^{(2)}}{f}, \quad v = 2(\ln f)_{xx}, \quad (27)$$

where $f, h^{(1)}$ and $h^{(2)}$ are given in the following Gram determinants:

$$f = \left| \delta_{ij} + \frac{1}{p_i + p_j^*} e^{\xi_i + \xi_j^*} \right|_{N \times N} , \qquad (28)$$

$$h^{(1)} = \left| \delta_{ij} + \left(-\frac{p_i - i\alpha_1}{p_j^* + i\alpha_1} \right) \frac{1}{p_i + p_j^*} e^{\xi_i + \xi_j^*} \right|_{N \times N},$$
(29)

$$h^{(2)} = \left| \delta_{ij} + \left(-\frac{p_i - i\alpha_2}{p_j^* + i\alpha_2} \right) \frac{1}{p_i + p_j^*} e^{\xi_i + \xi_j^*} \right|_{N \times N},$$
(30)

with

$$\theta_{1} = \alpha_{1}x - \alpha_{1}^{2}t + \beta_{1}(y), \quad \theta_{2} = \alpha_{2}x - \alpha_{2}^{2}t + \beta_{2}(y),$$

$$\xi_{i} = p_{i}x + ip_{i}^{2}t - \frac{1}{2}\left(\frac{\sigma_{1}k_{1}^{2}}{p_{i} - i\alpha_{1}} + \frac{\sigma_{2}k_{2}^{2}}{p_{i} - i\alpha_{2}}\right)y + \xi_{i0}, \quad (i = 1, 2, \dots, N),$$

1950390-6

where $\beta_1(y)$ and $\beta_2(y)$ are all arbitrary real functions of y, p_i and ξ_{i0} are complex constants, δ_{ij} is the Kronecker symbol.

3. Dynamics of the N-Dark-Dark Soliton Solutions

3.1. One dark-dark solitons

In this paper, the long wave component v always possesses bright soliton and we consider to construct dark solitons in the short wave components. In order to get one-dark-dark soliton for the two component coupled Maccari system (3)–(5), we choose N = 1 in the formulae (27)–(30). The concrete expressions in Gram type determinant forms can be given as

$$f_1 = 1 + \frac{1}{p_1 + p_1^*} e^{\xi_1 + \xi_1^*}, \qquad (31)$$

$$h_1^{(1)} = 1 - \frac{p_1 - i\alpha_1}{p_1^* + i\alpha_1} \frac{1}{p_1 + p_1^*} e^{\xi_1 + \xi_1^*} , \qquad (32)$$

$$h_1^{(2)} = 1 - \frac{p_1 - i\alpha_2}{p_1^* + i\alpha_2} \frac{1}{p_1 + p_1^*} e^{\xi_1 + \xi_1^*} , \qquad (33)$$

then substituting the above expressions into the dependent variable transformation (27), the one-dark-dark soliton solutions can read as

$$\phi^{(1)} = \frac{k_1}{2} e^{i\theta_1} \left[1 + H_1^{(1)} + (H_1^{(1)} - 1) \tanh\left(\frac{\xi_1 + \xi_1^* + \Theta_1}{2}\right) \right], \quad (34)$$

$$\phi^{(2)} = \frac{k_2}{2} e^{i\theta_2} \left[1 + H_1^{(2)} + (H_1^{(2)} - 1) \tanh\left(\frac{\xi_1 + \xi_1^* + \Theta_1}{2}\right) \right], \quad (35)$$

$$v = \frac{(p_1 + p_1^*)^2}{2} \operatorname{sech}^2\left(\frac{\xi_1 + \xi_1^* + \Theta_1}{2}\right),\tag{36}$$

where

$$e^{\Theta_1} = \frac{1}{p_1 + p_1^*} = \frac{1}{2m_1}, \quad H_1^{(1)} = -\frac{p_1 - i\alpha_1}{p_1^* + i\alpha_1} = -\frac{m_1 + i(n_1 - \alpha_1)}{m_1 - i(n_1 - \alpha_1)},$$
$$H_1^{(2)} = -\frac{p_1 - i\alpha_2}{p_1^* + i\alpha_2} = -\frac{m_1 + i(n_1 - \alpha_2)}{m_1 - i(n_1 - \alpha_2)}, \quad \xi_1 + \xi_1^* = 2m_1x - 4m_1n_1t_2$$
$$-\left(\frac{\sigma_1m_1k_1^2}{m_1^2 + (n_1 - \alpha_1)^2} + \frac{\sigma_2m_1k_2^2}{m_1^2 + (n_1 - \alpha_2)^2}\right)y + 2\xi_{10R},$$

here, $p_1 = m_1 + in_1$, $\xi_{10} = \xi_{10R} + i\xi_{10I}$ and m_1 , n_1 , ξ_{10R} , ξ_{10I} are all real constants. Choosing $m_1 > 0$, we can get the nonsingular one-dark-dark soliton solutions.

From Eqs. (34)–(36), we can directly calculate that the two short wave components' intensity functions $|\phi^{(1)}|$, $|\phi^{(2)}|$ and the long wave component's amplitude v move at velocity $2n_1$ along the x-direction, and at $\frac{-4n_1}{\frac{\sigma_1 k_1^2}{m_1^2 + (n_1 - \alpha_1)^2} + \frac{\sigma_2 k_2^2}{m_1^2 + (n_1 - \alpha_2)^2}}$ along

the y-direction. When $x, y \to \pm \infty$, we can derive that $|\phi^{(1)}| \to |k_1|, \phi^{(2)}| \to |k_2|$ and $v \to 0$.

The two complex constants $H_1^{(1)}$ and $H_1^{(2)}$ can be written as the following forms $H_1^{(1)} = e^{2i\psi_1^{(1)}}$ and $H_1^{(2)} = e^{2i\psi_1^{(2)}}$ $(-\frac{\pi}{2} \leq \psi_1^{(1)}, \psi_1^{(2)} \leq \frac{\pi}{2})$. With x and y varying from $-\infty$ to $+\infty$, the phase shifts of the two short wave components $\phi^{(1)}$ and $\phi^{(2)}$ are $2\psi_1^{(1)}$ and $2\psi_1^{(2)}$, respectively, but the phase shift of the long wave component v equals zero. Choosing $\xi_1 + \xi_1^* + \Theta_1 = 0$, we can directly calculate the intensities of the center for the three solitons as $|\phi^{(1)}|_{\text{center}} = |k_1| \cos \psi_1^{(1)}$, $|\phi^{(2)}|_{\text{center}} = |k_2| \cos \psi_1^{(2)}$ and $v_{\text{center}} = 2m_1^2$. For the two short wave components, it can be shown that the center intensities $|\phi^{(i)}|_{\text{center}} = |k_i| \cos \psi_1^{(i)}$ are smaller than the plane background $|k_i|$ and these two solitons are dark solitons. However, the soliton in the long wave component is bright. Additionally, the degrees of the darkness in the dark solitons in the two short wave components are controlled by the corresponding phase shifts $2\psi_1^{(1)}$ and $2\psi_1^{(2)}$.

According to the expressions of $H_1^{(1)}$ and $H_1^{(2)}$, we can discuss the degrees of the darkness for the two dark solitons in two different cases. (i) When $\alpha_1 = \alpha_2$, then $H_{1_{11}}^{(1)} = H_1^{(2)}$ and $\psi_1^{(1)} = \psi_1^{(2)}$. In this case, the two short

(i) When $\alpha_1 = \alpha_2$, then $H_1^{(1)} = H_1^{(2)}$ and $\psi_1^{(1)} = \psi_1^{(2)}$. In this case, the two short wave components $\phi^{(1)}$ and $\phi^{(2)}$ have the same degrees of darkness and they are proportional to each other. At this moment, the dark-dark solitons of the coupled Maccari system including two short waves are same as a scalar dark soliton in the Maccari system with one short wave. It can be seen as a degenerate case similar to the coupled NLS,¹⁸ YO,²³ Mel'nikov²⁴ equations. The degenerate case for the dark-dark solitons is shown in Fig. 1, then we can find that the valleys for the two dark solitons reach the zero plane and the long wave component possesses bright soliton.

(ii) When $\alpha_1 \neq \alpha_2$, then $H_1^{(1)} \neq H_1^{(2)}$ and $\psi_1^{(2)} \neq \psi_1^{(2)}$. At this point, the long wave component v still owns bright soliton and the valley of the dark soliton in



Fig. 1. (Color online) One dark-dark solitons in the degenerate case at the fixed time t = 0 with the parameters chosen by $k_1 = 1$, $k_2 = 2$, $m_1 = 1$, $n_1 = 1$, $\alpha_1 = 1$, $\alpha_2 = 1$, $\sigma_1 = 1$, $\sigma_2 = -1$, $\beta_1 = 0$, $\beta_2 = 0$, $\xi_{10R} = 0$.

1950390-8



Fig. 2. (Color online) One dark-dark solitons in the non-degenerate case at the fixed time t = 0 with the parameters chosen by $k_1 = 1$, $k_2 = 2$, $m_1 = 1$, $n_1 = 1$, $\alpha_1 = 1$, $\alpha_2 = 2$, $\sigma_1 = 1$, $\sigma_2 = -1$, $\beta_1 = 0$, $\beta_2 = 0$, $\xi_{10R} = 0$.

 $\phi^{(1)}$ component reaches the bottom, but the one in $\phi^{(2)}$ does not reach the bottom (which can also be called grey soliton), see Fig. 2. This is the non-degenerate case. From Fig. 2(a), it is shown that the two short wave components $\phi^{(1)}$ and $\phi^{(2)}$ have different degrees of darkness, and they are no longer proportional to each other. Furthermore, it is greatly different from the dark soliton in the Maccari system including one short wave component.

3.2. Two dark-dark solitons

Similarly, the two dark-dark solitons for the Maccari system (3)–(5) can be directly calculated with N = 2 in Eqs. (27)–(30). The two dark-dark solitons can be written as follows:

$$\phi^{(1)} = k_1 e^{i(\alpha_1 x - \alpha_1^2 + \beta_1(y))} \frac{h_2^{(1)}}{f_2}, \quad \phi^{(2)} = k_2 e^{i(\alpha_2 x - \alpha_2^2 + \beta_2(y))} \frac{h_2^{(2)}}{f_2}, \quad v = 2(\ln f_2)_{xx},$$
(37)

where

$$f_2 = 1 + e^{\xi_1 + \xi_1^* + \Theta_1} + e^{\xi_2 + \xi_2^* + \Theta_2} + \Lambda_{12} e^{\xi_1 + \xi_1^* + \xi_2 + \xi_2^* + \Theta_1 + \Theta_2},$$
(38)

$$h_{2}^{(1)} = 1 + H_{1}^{(1)}e^{\xi_{1} + \xi_{1}^{*} + \Theta_{1}} + H_{2}^{(1)}e^{\xi_{2} + \xi_{2}^{*} + \Theta_{2}} + H_{1}^{(1)}H_{2}^{(1)}\Lambda_{12}e^{\xi_{1} + \xi_{1}^{*} + \xi_{2} + \xi_{2}^{*} + \Theta_{1} + \Theta_{2}},$$
(39)

$$h_{2}^{(2)} = 1 + H_{1}^{(2)} e^{\xi_{1} + \xi_{1}^{*} + \Theta_{1}} + H_{2}^{(2)} e^{\xi_{2} + \xi_{2}^{*} + \Theta_{2}} + H_{1}^{(2)} H_{2}^{(2)} \Lambda_{12} e^{\xi_{1} + \xi_{1}^{*} + \xi_{2} + \xi_{2}^{*} + \Theta_{1} + \Theta_{2}},$$
(40)

with

$$e^{\Theta_j} = \frac{1}{p_j + p_j^*} = \frac{1}{2m_j}, \quad H_j^{(1)} = -\frac{p_j - i\alpha_1}{p_j^* + i\alpha_1} = -\frac{m_j + i(n_j - \alpha_1)}{m_j - i(n_j - \alpha_1)}, \tag{41}$$

$$H_{j}^{(2)} = -\frac{p_{j} - i\alpha_{2}}{p_{j}^{*} + i\alpha_{2}} = -\frac{m_{j} + i(n_{j} - \alpha_{2})}{m_{j} - i(n_{j} - \alpha_{2})},$$

$$\Lambda_{12} = \left|\frac{p_{1} - p_{2}}{p_{1} + p_{2}^{*}}\right|^{2} = \frac{(m_{1} - m_{2})^{2} + (n_{1} - n_{2})^{2}}{(m_{1} + m_{2})^{2} + (n_{1} - n_{2})^{2}},$$
(42)

$$\xi_j + \xi_j^* = 2m_j x - 4m_j n_j t - \left(\frac{\sigma_1 k_1^2 m_j}{m_j^2 + (n_j - \alpha_1)^2} + \frac{\sigma_2 k_2^2 m_j}{m_j^2 + (n_j - \alpha_2)^2}\right) y + 2\xi_{j0R}$$

$$\equiv L_{jx} x + L_{jt} t + L_{jy} y \quad (j = 1, 2), \qquad (43)$$

here, $p_j = m_j + in_j$, $\xi_{j0} = \xi_{j0R} + i\xi_{j0I}$ and m_j , n_j , α_1 , α_2 , ξ_{j0R} , ξ_{j0I} are real constants.

From the above expression of Λ_{12} , we find that the denominator becomes zero with the parameters chosen by $m_1 = -m_2$, $n_1 = n_2$. At this point, the two solitons interaction exists as Y-shape, which is called the resonant solitons.^{23,24} When Λ_{12} exists, we discuss the asymptotic analysis for the two solitons as follows.

(i) Before the interaction $(x, y \to -\infty)$: Soliton 1 s_1 $(\xi_{1R} \simeq 0, \xi_{2R} \to -\infty)$

$$\begin{split} \phi_1^{(1)-} &\simeq \frac{k_1}{2} e^{i\theta_1} \left[\left(1 + H_1^{(1)} + (H_1^{(1)} - 1) \right) \tanh\left(\frac{\xi_1 + \xi_1^* + \Theta_1}{2}\right) \right], \\ \phi_1^{(2)-} &\simeq \frac{k_2}{2} e^{i\theta_2} \left[\left(1 + H_1^{(2)} + (H_1^{(2)} - 1) \right) \tanh\left(\frac{\xi_1 + \xi_1^* + \Theta_1}{2}\right) \right], \\ v_1^- &\simeq \frac{(p_1 + p_1^*)^2}{2} \operatorname{sech}^2 \left(\frac{\xi_1 + \xi_1^* + \Theta_1}{2}\right), \end{split}$$

Soliton 2 $s_2 \ (\xi_{2R} \simeq 0, \xi_{1R} \to +\infty)$

$$\begin{split} \phi_2^{(1)-} &\simeq \frac{k_1 H_1^{(1)}}{2} e^{i\theta_1} \left[(1+H_2^{(1)}) + (H_2^{(1)}-1) \tanh\left(\frac{\xi_2 + \xi_2^* + \Theta_2 + \ln\Lambda_{12}}{2}\right) \right], \\ \phi_2^{(2)-} &\simeq \frac{k_2 H_1^{(2)}}{2} e^{i\theta_2} \left[(1+H_2^{(2)}) + (H_2^{(2)}-1) \tanh\left(\frac{\xi_2 + \xi_2^* + \Theta_2 + \ln\Lambda_{12}}{2}\right) \right], \\ v_2^- &\simeq \frac{(p_2 + p_2^*)^2}{2} \operatorname{sech}^2 \left(\frac{\xi_2 + \xi_2^* + \Theta_2 + \ln\Lambda_{12}}{2}\right), \end{split}$$

(ii) After the interaction $(x, y \to +\infty)$: Soliton 1 s_1 $(\xi_{1R} \simeq 0, \xi_{2R} \to +\infty)$

$$\phi_1^{(1)+} \simeq \frac{k_1 H_2^{(1)}}{2} e^{i\theta_1} \left[(1+H_1^{(1)}) + (H_1^{(1)}-1) \tanh\left(\frac{\xi_1 + \xi_1^* + \Theta_1 + \ln\Lambda_{12}}{2}\right) \right].$$

Multi-dark soliton solutions for the (2+1)-dimensional multi-component Maccari system

$$\begin{split} \phi_1^{(2)+} &\simeq \frac{k_2 H_2^{(2)}}{2} e^{i\theta_2} \left[(1+H_1^{(2)}) + (H_1^{(2)}-1) \tanh\left(\frac{\xi_1 + \xi_1^* + \Theta_1 + \ln\Lambda_{12}}{2}\right) \right],\\ v_1^+ &\simeq \frac{(p_1 + p_1^*)^2}{2} \operatorname{sech}^2\left(\frac{\xi_1 + \xi_1^* + \Theta_1 + \ln\Lambda_{12}}{2}\right), \end{split}$$

Soliton 2 $s_2 \ (\xi_{2R} \simeq 0, \xi_{1R} \to -\infty)$

$$\begin{split} \phi_2^{(1)+} &\simeq \frac{k_1}{2} e^{i\theta_1} \left[(1+H_2^{(1)}) + (H_2^{(1)}-1) \tanh\left(\frac{\xi_2 + \xi_2^* + \Theta_2}{2}\right) \right], \\ \phi_2^{(2)+} &\simeq \frac{k_2}{2} e^{i\theta_2} \left[(1+H_2^{(2)}) + (H_2^{(2)}-1) \tanh\left(\frac{\xi_2 + \xi_2^* + \Theta_2}{2}\right) \right], \\ v_2^+ &\simeq \frac{(p_2 + p_2^*)^2}{2} \operatorname{sech}^2 \left(\frac{\xi_2 + \xi_2^* + \Theta_2}{2}\right), \end{split}$$

where $\xi_j = \xi_{jR} + i\xi_{jI}(j = 1, 2)$, $\phi_j^{(1)-}$, $\phi_j^{(2)-}$ denotes the two solitons for the two short wave components before collision, and v_j^- are the two solitons for long wave component before collision. Conversely, $\phi_j^{(1)+}$, $\phi_j^{(2)+}$ and v_j^+ stand for the two solitons in the corresponding components after collision.

From the above expressions of asymptotic analysis, we can directly calculate that

$$\begin{aligned} A_1^{(1)-} &= A_1^{(1)+} = |k_1| \left(1 - \frac{\sqrt{\frac{1}{3}m_1^2 + (n_1 - \alpha_1)^2}}{m_1^2 + (n_1 - \alpha_1)^2} \right), \\ A_1^{(2)-} &= A_1^{(2)+} = |k_2| \left(1 - \frac{\sqrt{\frac{1}{3}m_1^2 + (n_1 - \alpha_1)^2}}{m_1^2 + (n_1 - \alpha_1)^2} \right), \\ A_2^{(1)-} &= A_2^{(1)+} = |k_1| \left(1 - \frac{\sqrt{\frac{1}{3}m_2^2 + (n_2 - \alpha_1)^2}}{m_2^2 + (n_2 - \alpha_1)^2} \right), \\ A_2^{(2)-} &= A_2^{(2)+} = |k_2| \left(1 - \frac{\sqrt{\frac{1}{3}m_2^2 + (n_2 - \alpha_2)^2}}{m_2^2 + (n_2 - \alpha_2)^2} \right), \\ A_v^{(1)-} &= A_v^{(1)+} = \frac{(p_1 + p_1^*)^2}{2} = 2m_1^2, \quad A_v^{(2)-} = A_v^{(2)+} = \frac{(p_2 + p_2^*)^2}{2} = 2m_2^2, \end{aligned}$$
(45)

where $A_1^{(j)-}$, $A_1^{(j)+}$ (j = 1, 2) denote the amplitude of the soliton j (s_j) in the short wave component $\phi^{(1)}$ before and after collision, respectively. Analogously, $A_2^{(j)-}$, $A_2^{(j)+}$ are the amplitudes of the soliton j in the short wave component $\phi^{(2)}$ before and after collision. Additionally, the amplitudes of the corresponding soliton s_j in the long wave component v can be written as $A_v^{(j)-}$ (before collision) and $A_v^{(j)+}$ (after collision). From the asymptotic analysis, we can make a conclusion that the amplitude and velocity of each soliton remain unchanged during the interaction



Fig. 3. (Color online) Two dark-dark solitons at the fixed time t = 0 with the parameters chosen by $k_1 = 1, k_2 = 2, m_1 = 1, n_1 = 2, m_2 = 2, n_2 = \frac{1}{3}, \alpha_1 = 1, \alpha_2 = 2, \sigma_1 = 1, \sigma_2 = -1, \beta_1 = 0, \beta_2 = 0, \xi_{10R} = 0, \xi_{20R} = 0.$

except for some phase shifts, and the collisions in the three components are all elastic.

From Eq. (38), we can easily get a series of nonsingular two dark-dark soliton solutions with the parameters chosen by $m_j > 0$ (j = 1, 2). Meantime, the collision of two dark-dark solitons is exhibited in Fig. 3. It is shown that the two solitons interact with each other without any change of amplitude, darkness and velocity in the three components after collision. There is no energy exchange between the two short wave components or two dark solitons in the same component after interaction. This kind of transmission of energy of dark-dark soliton solutions occurs in all possible cases of nonlinearity coefficients σ_1 and σ_2 (all positive, all negative and mixed).

For the two bright solitons in the long wave component v, we can differentiate them into two types (O-type and P-type) according to asymptotic analysis. From Eq. (45), we can find that the amplitudes of the two solitons s_1 and s_2 for the long wave component in the asymptotic analysis are $2m_1^2$ and $2m_2^2$, respectively. According to the classifications of the soliton solutions in the Kadomtsev–Petviashvili (KP) equation,^{44,45} the two bright solitons in v can be classified as O-type ("O" for original) and P-type ("P" for physical) soliton interactions.

(i) When $m_1m_2 < 0$, then $\Lambda_{12} > 1$, this type is called O-type soliton interaction. The two asymptotic soliton amplitudes $2m_1^2$ and $2m_2^2$ for the long wave component v can be equivalent when $m_1 = -m_2$. It is found that the maximum of v (interaction peak) is always bigger than the sum of the asymptotic soliton $(s_1 \text{ and } s_2)$ amplitudes.

(ii) When $m_1m_2 > 0$, then $0 < \Lambda_{12} < 1$, this type is called P-type soliton interaction. The following equality that $2m_1^2 = 2m_2^2$ $(m_1 \neq m_2)$ is not possible in this case. It is shown that the maximum of v (interaction peak) is always smaller than the sum of the asymptotic soliton amplitudes. The two bright solitons in Fig. 3(c) are just P-type solitons interaction.

It should be noted that the two different types of soliton interactions do not depend on the imaginary parts of p_1 and p_2 (namely n_1 and n_2). The abovementioned Y-shape soliton solution can be constructed through using a limiting process $n_1 \rightarrow n_2$ in the O-type two solitons. Similar with the cases in one- and two dark-dark solitons, the higher-order dark-dark solitons for the Maccari system (3)–(5) can also be discussed and we omit them here.

4. Soliton Bound States

In the following contents, we will consider the soliton bound states in the three components $\phi^{(1)}$, $\phi^{(2)}$ and v. In order to construct the bound states for the Maccari system (3)–(5), the solitons of the short and long wave components should possess the same velocities along the x- and y-direction. From the concrete expressions (37)–(40) of two dark-dark solitons, we can directly calculate the two velocities V_{jx} (along the x-direction) and V_{jy} (along the y-direction) as follows (j = 1, 2 denote the two different solitons)

$$V_{jx} = 2n_j, \quad V_{j,y} = -\frac{4n_j}{\frac{\sigma_1 k_1^2}{m_j^2 + (n_j - \alpha_1)^2} + \frac{\sigma_2 k_2^2}{m_j^2 + (n_j - \alpha_2)^2}} \quad (j = 1, 2).$$
(46)

When $V_{jx} = V_{jy} = 0$, we can get the stationary dark-dark soliton bound states. From Eq. (46), it only requires that $n_1 = n_2 = 0$. There are no other constraints for the nonlinear coefficients, then all values of σ_1, σ_2 (all positive, all negative and mixed) can satisfy this condition. However, the relative constraints only exist when the nonlinearity coefficients take opposite signs in the coupled NLS equations¹⁸ and YO equations.²³ The higher-order stationary dark-dark soliton bound states only require that $n_1 = n_2 = \cdots = n_i (3 \le i \le N)$, and we can make a conclusion that the bound states can exist up to arbitrary order in the stationary case.

According to the relationship between the coefficients L_{jx} and L_{jy} in Eq. (43), the stationary bound states can be classified as two types — oblique and parallel. (i) If $\frac{L_{1y}}{L_{1x}} \neq \frac{L_{2y}}{L_{2x}}$, namely $\frac{\sigma_1 k_1^2}{m_1^2 + \alpha_1^2} + \frac{\sigma_2 k_2^2}{m_1^2 + \alpha_2^2} \neq \frac{\sigma_1 k_1^2}{m_2^2 + \alpha_1^2} + \frac{\sigma_2 k_2^2}{m_2^2 + \alpha_2^2}$, this is the oblique stationary bound state, see Fig. 4. Here, it is shown that the coefficients of the independent variable t are all zero in the expressions of two dark-dark solitons, and these two solitons do not propagate along time t. Moreover, the oblique



Fig. 4. (Color online) The stationary two dark-dark soliton bound states in the oblique case with the parameters chosen by $k_1 = 1$, $k_2 = 2$, $m_1 = 1$, $n_1 = 0$, $m_2 = \sqrt{3}$, $n_2 = 0$, $\alpha_1 = 1$, $\alpha_2 = \frac{1}{3}$, $\sigma_1 = 1$, $\sigma_2 = 1$, $\beta_1 = 0$, $\beta_2 = 0$, $\xi_{10R} = 0$, $\xi_{20R} = 0$.



Fig. 5. (Color online) The stationary two dark-dark soliton bound states in the parallel case with the parameters chosen by $k_1 = 1$, $k_2 = 2$, $m_1 = 1$, $n_1 = 0$, $m_2 = \frac{1}{33}\sqrt{671}$, $n_2 = 0$, $\alpha_1 = \frac{1}{3}$, $\alpha_2 = 1$, $\sigma_1 = 1$, $\sigma_2 = -1$, $\beta_1 = 0$, $\beta_2 = 0$, $\xi_{10R} = 0$, $\xi_{20R} = 0$.

stationary bound state can be generated for all possible combinations of nonlinearity coefficients σ_1 , σ_2 consisting of positive, negative and mixed cases.

(ii) If $\frac{L_{1y}}{L_{1x}} = \frac{L_{2y}}{L_{2x}}$, namely $\frac{\sigma_1 k_1^2}{m_1^2 + \alpha_1^2} + \frac{\sigma_2 k_2^2}{m_1^2 + \alpha_2^2} = \frac{\sigma_1 k_1^2}{m_2^2 + \alpha_1^2} + \frac{\sigma_2 k_2^2}{m_2^2 + \alpha_2^2}$, this is the parallel stationary bound state, see Fig. 5. From Eq. (38), one of the constraint conditions for constructing the nonsingular two dark-dark solitons is that $m_1 > 0, m_2 > 0$. Given the following function $f(x) = \frac{\sigma_1 k_1^2}{x^2 + \alpha_1^2} + \frac{\sigma_1 k_1^2}{x^2 + \alpha_2^2}$ (x > 0), and the parallel stationary bound state exists if and only if $f(m_1) = f(m_2)$ by choosing appropriate values of $\sigma_1, \sigma_2, k_1, k_2, \alpha_1, \alpha_2$. Besides, the function f(x) is monotonous in the interval x > 0 when the nonlinearity coefficients σ_1, σ_2 take the same signs (all positive or all negative). Considering the above condition $m_1 > 0, m_2 > 0$, it is not possible that $f(m_1) = f(m_2)$ when $m_1 \neq m_2, \sigma_1 \sigma_2 > 0$. Based on the above facts, we find that the parallel stationary is only possible when nonlinearity coefficients σ_1, σ_2 take opposite signs.

When $V_{jx} = V_{jy} \neq 0$, the moving two dark-dark soliton bound states can be constructed. From Eq. (46), the following restricted conditions for constructing moving bound states can be written as

$$V_{jx}: n_1 = n_2 \neq 0,$$

$$V_{jy}: \frac{\sigma_1 k_1^2}{m_1^2 + (n_1 - \alpha_1)^2} + \frac{\sigma_2 k_2^2}{m_1^2 + (n_1 - \alpha_2)^2} \qquad (47)$$

$$= \frac{\sigma_1 k_1^2}{m_2^2 + (n_1 - \alpha_1)^2} + \frac{\sigma_2 k_2^2}{m_2^2 + (n_1 - \alpha_2)^2}.$$

Similar with the case in parallel stationary bound state, we can also construct the function $g(x) = \frac{\sigma_1 k_1^2}{x^2 + (n_1 - \alpha_1)^2} + \frac{\sigma_2 k_2^2}{x^2 + (n_1 - \alpha_2)^2}$ (x > 0). The function g(x) is monotonous in the interval x > 0 with fixed parameters $k_1, k_2, n_1, \alpha_1, \alpha_2$ when nonlinearity coefficients σ_1, σ_2 take the same signs. Considering the nonsingular condition $m_1 > 0, m_2 > 0$, it is not possible that $g(m_1) = g(m_2)$ ($m_1 \neq m_2$) if $\sigma_2 \sigma_2 > 0$. It is shown that the moving bound states are only possible when nonlinearity coefficients σ_1, σ_2 take opposite signs.



Fig. 6. (Color online) (a), (b): The moving two dark-dark soliton bound states at time t = -8 in the two short wave components; (c) the profile of two bright-bright soliton bound states with y = 10 at t = -8 in the long wave component.



Fig. 7. (Color online) (a), (b): The moving two dark-dark soliton bound states at time t = 0 in the two short wave components; (c) the profile of two bright-bright soliton bound states with y = 10 at t = 0 in the long wave component.

Assuming that the moving bound states can exist up to three dark solitons, and we have $n_1 = n_2 = n_3$, $g(m_1) = g(m_2) = g(m_3)(m_1, m_2, m_3 > 0)$. From the expression of g(x), we can find that g(x) is the equation with one independent variable x whose degree is four, and the equation g(x) = c has two positive roots at most. Based on the above facts, it is not possible that $g(m_1) = g(m_2) = g(m_3)$ with $m_1, m_2, m_3 > 0$. A conclusion can be made that only two-soliton bound state exists in the moving case. Additionally, the figures of the moving two dark-dark solitons are shown in Figs. 6–8 at three different times with the parameters chosen by

$$\begin{aligned} k_1 &= 1, \quad k_2 = 2, \quad m_1 = 1, \quad n_1 = 1, \quad m_2 = \frac{5}{52}\sqrt{39}, \quad n_2 = 1, \quad \alpha_1 = 1, \\ \alpha_2 &= \frac{1}{4}, \quad \sigma_1 = 1, \quad \sigma_2 = -1, \quad \beta_1 = 0, \quad \beta_2 = 0, \quad \xi_{10R} = 0, \quad \xi_{20R} = 0. \end{aligned}$$

We find that it is not easy to observe the moving two bright-bright solitons in the long wave component v, and we give the corresponding two-dimensional diagrams at y = 10 in Figs. 6(c)-8(c).

For the two-soliton bound states (stationary and moving cases), it is interesting that the two short wave components $\phi^{(1)}$ and $\phi^{(2)}$ have some phase shifts but the long wave component v undergoes no phase shift with x and y varying from $-\infty$



Fig. 8. (Color online) (a), (b): The moving two dark-dark soliton bound states at time t = 8 in the two short wave components; (c) the profile of two bright-bright soliton bound states with y = 10 at t = 8 in the long wave component.

to $+\infty$. Additionally, the sum of the phase shifts of each short wave component are equal to the sum of the individual ones of the two dark solitons. Whereas the phase shift of the long wave component is still zero. Setting that $H_j^{(1)} = e^{2i\psi_j^{(1)}}$, $H_j^{(2)} = e^{i\psi_j^{(2)}}$, then the phase shifts of the corresponding components can be written as $\phi_{\text{phase shift}}^{(1)} = 2\psi_1^{(1)} + 2\psi_2^{(1)}$, $\phi_{\text{phase shift}}^{(2)} = 2\psi_1^{(2)} + 2\psi_2^{(2)}$ and $v_{\text{phase shift}} = 0$.

5. N-Dark Soliton Solutions for the Multi-Component Generalization

If the isolated waves localized in small parts of space are more than two, the Maccari system (3) and (5) should be generalized to multi-component case. In fact, the previous results for the two component case can be extended to the N-component Maccari system (1) and (2). Furthermore, the multi-dark soliton solutions for the N-component Maccari system can be generated from the reduction of single KP hierarchy consisting of N copies of shifted singular points. Similar with the processes in the two component coupled Maccari system (3) and (5), we can directly calculate the N-dark soliton solutions for the N-component Maccari system (1) and (2). Here, we only give the main results and omit some complicated calculation procedures.

The N-component Maccari system (1) and (2) consisting of N short wave components and single long wave component can be transformed into the following bilinear forms:

$$[D_x^2 + i(D_t + 2\alpha_l D_x)]h^{(l)} \cdot f = 0 \quad (1 \le l \le N),$$
(48)

$$D_x D_y f \cdot f = \sum_{l=1}^N \sigma_l k_l^2 (h^{(l)} h^{(l)*} - f^2), \qquad (49)$$

through these dependent variable transformations

$$\phi^{(l)} = k_l e^{i\theta_l} \frac{h^{(l)}}{f} \quad (1 \le l \le N), \quad v = 2(\ln f)_{xx}, \tag{50}$$

where $h^{(l)}(1 \le l \le N)$ are the complex functions of x, y, t, f is the real function of x, y, t. Besides, $\theta_l = \alpha_l x - \alpha_l^2 t + \beta_l(y), k_l, \alpha_l$ are arbitrary real constants and $\beta_l(y)$ are the arbitrary functions of y.

Similar to the reductions in Sec. 2, one can construct the N-dark soliton solutions in Gram type determinant forms for the multi-component Maccari system (1) and (2) as follows:

$$\phi^{(l)} = k_l e^{i\theta_l} \frac{h^{(l)}}{f}, \quad v = 2(\ln f)_{xx},$$
(51)

$$f = \left| \delta_{ij} + \frac{1}{p_i + p_j^*} e^{\xi_i + \xi_j^*} \right|_{N \times N},$$
 (52)

$$h^{(l)} = \left| \delta_{ij} - \frac{p_i - i\alpha_l}{p_j^* + i\alpha_l} \frac{1}{p_i + p_j^*} \right|_{N \times N} \quad (1 \le l \le N),$$
(53)

where $\xi_i = p_i x + i p_i^2 - \frac{1}{2} \sum_{l=1}^N \frac{\sigma_l k_l^2}{p_i - i \alpha_l} y + \xi_{i0}$, $\theta_l = \alpha_l x - \alpha_l^2 t + \beta_l(y)$, $\beta_l(y)$ are the arbitrary functions of y, p_i , ξ_{i0} are complex constants and k_l , α_l are real constants, δ_{ij} is the Kronecker symbol.

6. Conclusions

Utilizing reduction of the KP hierarchy, the N-dark-dark soliton solutions for the two component coupled Maccari system (3)–(5) consisting of two short wave components and one long wave component are given in Gram determinant forms. There exist some connections between the researched Maccari system and some equations in the KP hierarchy. Here, the related equations are the two-dimensional Toda lattice (Eqs. (11) and (13)) in KP hierarchy and the lowest-degree bilinear equation (Eqs. (12) and (14)) in the first modified KP hierarchy. Based on the above facts, the N-dark solitons for the Maccari system (3)–(5) can be constructed by the complex conjugate reductions and the independent variable transformations in the KP hierarchy. Similar to the N-dark-dark soliton in the two component case, the N dark solitons for the N-component Maccari system including N short wave components and one long wave component can also be generated. This kind of dark soliton in two-dimensional multi-component system has also been reported for the YO system²³ and Mel'nikov system.²⁴

In this paper, the dynamics of the dark solitons for the two component coupled Maccari system are discussed in detail. For one-dark-dark solitons, the degrees of the darkness can be controlled by the parameter α_j (j = 1, 2) and they are classified as two types (degenerate and non-degenerate cases). From the asymptotic analysis, it is shown the amplitude and velocity of each soliton in the two-soliton solutions remains unchanged during the interaction, then the collisions in the three components are all elastic. The long wave component v always possesses bright soliton and the two bright soliton interactions are O-type and P-type. Additionally, the two dark-dark solitons bound states are discussed in stationary and moving cases. For the stationary bound states, the bound states can exist up to arbitrary order. However, only two-soliton bound state exists in the moving case. There are

two types (oblique and parallel) in the stationary bound states. The oblique case can be generated for all possible combinations of nonlinearity coefficients. Nevertheless, the parallel case is only possible when nonlinearity coefficients take opposite signs. We expect that the N-dark soliton obtained in this paper will be verified and controlled in the physical experiments in the future.

Acknowledgments

This paper is supported by ECNU scholarship program for graduate students short-term overseas research scholarship. The first and second authors' work are supported by the National Natural Science Foundation of China (Nos. 11675054 and 11435005), Global Change Research Program of China (No. 2015CB953904), Shanghai Collaborative Innovation Center of Trustworthy Software for Internet of Things (No. ZF1213), Science and Technology Commission of Shanghai Municipality (No. 18dz2271000). The third author's work is partially supported by the President's Endowed Professorship program of the University of Texas system.

References

- A. Hasegawa and Y. Kodama, Solitons in Optical Communication (Oxford University Press, New York, 1995).
- 2. D. Mihalache, Rom. Rep. Phys. 69 (2017) 403.
- 3. P. G. Kevrekidis and D. J. Frantzeskakis, Rev. Phys. 1 (2016) 140.
- 4. Y. H. Qin, L. C. Zhao, Z. Y. Yang and W. L. Yang, Chaos 28 (2018) 013111.
- 5. V. B. Matveev and A. O. Smirnov, J. Math. Phys. 59 (2018) 091419.
- 6. L. M. Ling, L. C. Zhao and B. L. Guo, Nonlinearity 28 (2015) 3243.
- 7. J. C. Chen, B. F. Feng, K. Maruno and Y. Ohta, Stud. Appl. Math. 141 (2018) 145.
- 8. X. Wang, C. Liu and L. Wang, Chaos 27 (2017) 093106.
- 9. L. M. Ling, B. L. Guo and L. C. Zhao, Phys. Rev. E 89 (2014) 041201(R).
- 10. D. J. Kaup and A. C. Newell, J. Math. Phys. 19 (1978) 798.
- 11. Z. J. Qiao, J. Math. Phys. 35 (1994) 2971.
- S. H. Chen, Y. L. Ye, J. M. Soto-Crespo, P. Grelu and F. Baronio, *Phys. Rev. Lett.* 121 (2018) 104101.
- 13. G. Q. Zhang and Z. Y. Yan, Commun. Nonlinear Sci. Numer. Simul. 62 (2018) 117.
- 14. X. E. Zhang, T. Xu and Y. Chen, Nonlinear Dynam. 94 (2018) 2841.
- 15. G. Mu, Z. Y. Qin and R. Grimshaw, SIAM J. Appl. Math. 75 (2015) 1.
- 16. T. Xu and Y. Chen, Nonlinear Dynam. 92 (2018) 2133.
- 17. L. M. Ling, L. C. Zhao and B. L. Guo, Nonlinearity 28 (2015) 3243.
- 18. Y. Ohta, D. S. Wang and J. K. Yang, Stud. Appl. Math. 127 (2011) 345.
- 19. C. Kalla, J. Phys. A: Math. Theor. 44 (2011) 335210.
- A. de O. Assuncao, H. Blas and M. J. F. da Silva, J. Phys. A: Math. Theor. 45 (2012) 085205.
- 21. H. Q. Zhang and S. S. Yuan, Nonlinear Dynam. 89 (2017) 531.
- 22. G. Q. Zhang, Z. Y. Yan and X. Y. Wen, Chaos. 27 (2017) 083110.
- 23. J. C. Chen, Y. Chen, B. F. Feng and K. Maruno, J. Phys. Soc. Jpn. 84 (2015) 034002.
- 24. Z. Han, Y. Chen and J. C. Chen, J. Phys. Soc. Jpn. 86 (2017) 074005.
- 25. M. Jimbo and T. Miwa, Publ. RIMS Kyoto Univ. 19 (1983) 943.
- 26. B. F. Feng, J. Phys. A: Math. Theor. 47 (2014) 355203.

- 27. J. C. Chen, Y. Chen, B. F. Feng and K. Maruno, J. Phys. Soc. Jpn. 84 (2015) 074001.
- 28. Z. Han and Y. Chen, Chinese Phys. Lett. 7 (2017) 070202.
- 29. Y. Ohta and J. K. Yang, Proc. R. Soc. A 468 (2012) 1716.
- J. C. Chen, Y. Chen, B. F. Feng, K. Maruno and Y. Ohta, J. Phys. Soc. Jpn. 87 (2018) 094007.
- 31. X. E. Zhang and Y. Chen, Nonlinear Dynam. 93 (2018) 2169.
- 32. J. C. Chen, Y. Chen, B. F. Feng and K. Marunod, (2017) arXiv:1712.00945v1 [nlin.SI].
- 33. A. Maccari, J. Math. Phys. 38 (1997) 4151.
- 34. A. B. Shabat and V. E. Zakharov, Sov. Phys. JETP 34 (1972) 62.
- 35. A. Fokas, *Inverse Probl.* **10** (1994) L19.
- A. D. D. Craik, Wave Interactions and Fluid Flows (Cambridge University Press, Cambridge, 1985).
- 37. F. Yuan, J. G. Rao, K. Porsezian, D. Mihalache and J. S. He, *Rom. J. Phys.* **61** (2016) 378.
- 38. S. Y. Lou, J. Lin and X. Y. Tang, Eur. Phys. J. B 22 (2001) 473.
- 39. W. H. Huang, Y. L. Liu and Z. Y. Ma, Commun. Theor. Phys. 47 (2007) 397.
- 40. J. F. Zhang, Commun. Theor. Phys. 35 (2001) 267.
- A. Uthayakumar, K. Nakkeeran and K. Porsezian, *Chaos Soliton. Fract.* 10 (1999) 1513.
- 42. Z. Han and Y. Chen, Z. Naturforsch. A 72 (2017) 745.
- 43. E. Date, M. Jimbo and T. Miwa, J. Phys. Soc. Jpn. 51 (1982) 4125.
- 44. S. Chakravarty, T. Lewkow and K. Maruno, Appl. Anal. 89 (2010) 529.
- 45. S. Chakravarty and Y. Kodama, J. Phys. A 41 (2008) 275209.