PAPER

High-order rational solutions and resonance solutions for a (3+1)–dimensional Kudryashov-Sinelshchikov equation^{*}

To cite this article: Yun-Fei Yue et al 2021 Chinese Phys. B 30 010202

View the article online for updates and enhancements.



You may also like

- The Ising model and special geometries S Boukraa, S Hassani and J-M Maillard
- <u>General Higher-Order Breather and Hybrid</u> <u>Solutions of the Fokas System</u> Ting-Ting Chen, , Peng-Yan Hu et al.
- <u>Resonant multi-soliton and multiple rogue</u> wave solutions of (3+1)-dimensional <u>Kudryashov-Sinelshchikov equation</u> Yueyang Feng and Sudao Bilige

High-order rational solutions and resonance solutions for a (3+1)-dimensional Kudryashov–Sinelshchikov equation*

Yun-Fei Yue(岳云飞)¹, Jin Lin(林机)², and Yong Chen(陈勇)^{1,2,3,†}

¹ School of Mathematical Sciences, Shanghai Key Laboratory of PMMP, Shanghai Key Laboratory of Trustworthy Computing,

East China Normal University, Shanghai 200062, China

² Department of Physics, Zhejiang Normal University, Jinhua 321004, China

³College of Mathematics and Systems Science, Shandong University of Science and Technology, Qingdao 266590, China

(Received 10 August 2020; accepted manuscript online 1 September 2020)

We mainly investigate the rational solutions and *N*-wave resonance solutions for the (3+1)-dimensional Kudryashov– Sinelshchikov equation, which could be used to describe the liquid containing gas bubbles. With appropriate transformations, two kinds of bilinear forms are derived. Employing the two bilinear equations, dynamical behaviors of nine district solutions for this equation are discussed in detail, including bright rogue wave-type solution, dark rogue wave-type solution, bright W-shaped solution, dark W-shaped rational solution, generalized rational solution and bright-fusion, darkfusion, bright-fission, and dark-fission resonance solutions. In addition, the generalized rational solutions, which depending on two arbitrary parameters, have an interesting structure: splitting from two peaks into three peaks.

Keywords: rational solution, N-wave resonance solution, Hirota bilinear method, Kudryashov–Sinelshchikov equation

PACS: 02.30.IK, 05.45.Yv, 04.20.Jb

1. Introduction

It is well known that the research on wave propagation is one of the important problems in bubble liquid. The dynamics of pressure waves in the gas–liquid mixtures are characterized by the Burgers–Korteweg–de Vries (BKdV) equation and the Korteweg–de Vries (KdV) equation.^[1] The (3+1)-dimensional Kudryashov–Sinelshchikov (KS) equation with the effects of liquid viscosity and heat transfer on the phase interface is

$$\left(\rho_{t} + (1 + \frac{a_{1}}{a})\rho\rho_{x} + \frac{b}{2a}\rho_{xxx} - \frac{c}{2a}\right)_{x} + \frac{d^{2}}{2}\rho_{yy} + \frac{e^{2}}{2}\rho_{zz} = 0.$$
 (1)

Here the long waves propagate along the *x* axis and allow for horizontal development in the *y* and *z* directions. The solution $\rho = \rho(t, x, y, z)$ represents the density in the bubble–liquid mixture, the parameters *a*, *a*₁, *b*, and *c* are constants relying on the radius of bubbles and the pressure of the gas in the unperturbed state, the kinematic viscosity and the surface tension of the liquid, the polytrophic exponent, and the volume of gas in the per unit mass of the bubble–liquid mixture, *etc*. The parameters *d* and *e* denote the perturbations in the *y* and *z* directions.

With the scaling transformation

$$t \to \frac{2a}{b\beta^2}t, \ x \to \frac{1}{\beta}x, \ y \to \frac{d}{\beta}\sqrt{\frac{a}{b\beta\gamma}}y,$$

DOI: 10.1088/1674-1056/abb3f3

$$z \to \frac{e}{\beta} \sqrt{\frac{a}{b\beta\delta}} z, \ \rho \to \frac{\alpha\beta b}{2(a+a_1)} u,$$
 (2)

the three-dimensional KS equation (1) can be converted into the following form:

$$u_{xt} + \alpha (uu_x)_x + \beta u_{xxxx} - \chi u_{xxx} + \gamma u_{yy} + \delta u_{zz} = 0, \quad (3)$$

where $\chi = c/b$. It is noted that this equation has very important physical implications, and can be reduced to some classical physical models by choosing suitable parameters in the following cases:

(i) The two-dimensional BKdV equation^[2] ($\delta = 0$);

(ii) The KdV equation^[3] ($\alpha = 6, \beta = 1, \chi = \gamma = \delta = 0$);

(iii) The KP equation^[4] ($\alpha = 6$, $\beta = 1$, $\chi = \delta = 0$, $\gamma = \pm 1$);

(iv) The one-dimensional KS^[5] or BKdV equation^[6] $(\beta = 1, \gamma = \delta = 0);$

(v) The two-dimensional Burgers-KP (BKP) equation^[6] ($\alpha = 1, \beta = \delta = 0$);

(vi) The three-dimensional KP equation^[7] ($\alpha = 6, \beta = 1, \chi = 0, \delta = \gamma = \pm 3$).

In recent years, several attempts have been made to construct explicit solutions for some nonlinear systems.^[9–15] It is a difficult and complicate but very important and meaningful work to find the nonlinear wave solutions and interaction solutions for nonlinear systems. By solving the nonlinear mathematical physics model, many kinds of nonlinear physical phe-

Corresponding aution. E-mail. yenen@sel.eenu.edu.en

^{*}Project supported by the National Natural Science Foundation of China (Grant No. 11675054), the Future Scientist/Outstanding Scholar Training Program of East China Normal University (Grant No. WLKXJ2019-004), the Fund from Shanghai Collaborative Innovation Center of Trustworthy Software for Internet of Things (Grant No. ZF1213), and the Project from the Science and Technology Commission of Shanghai Municipality, China (Grant No. 18dz2271000). [†]Corresponding author. E-mail: ychen@sei.ecnu.edu.cn

^{© 2021} Chinese Physical Society and IOP Publishing Ltd

nomena can be effectively described in nature. With the deepening of the nonlinear system research, a series of effective methods have formed for solving nonlinear partial differential equations, such as Hirota method,^[16] Darboux transformation method,^[17] KP reduction method,^[18] nonlocal symmetry method,^[19] and so on.

Consideration here is the issue of rational solutions for the (3+1)-dimensional KS equation only with the highest derivative terms of each variables, namely

$$u_{xt} + \alpha (uu_x)_x + \beta u_{xxxx} + \gamma u_{yy} + \delta u_{zz} = 0, \qquad (4)$$

where α , β , γ , and δ are real constants. It can be used to describe the liquid containing gas bubbles without considering its viscosity. To our knowledge, localized wave solutions, periodic wave solutions and interaction solutions of the (3+1)-dimensional KS equation (4) have been obtained in Refs. [20,21] Moreover, bright soliton, resonant multiple wave, lump-stripe, breather and rogue wave solutions have been derived for the (3+1)-dimensional KS equation with the scaling u(t,x,y,z) = u(t - hx, x, y, z).^[22–25] Motivated from the work by Clarkson and Dowie in a study of rational solutions to the Boussinesq equation,^[26] the aim of the present paper is to construct rogue wave-type rational solution, W-shaped rational solution, multiple basic rogue wave solution, and *N*-wave resonance solution of the (3+1)-dimensional KS equation (4), which have not been reported before.

In the present paper, a total of nine district solutions of Eq. (4) will be obtained by the Hirota bilinear method,^[16] including bright rogue wave-type rational solution, dark rogue wave-type rational solution, bright W-shaped rational solution, dark W-shaped rational solution, generalized rational solution, bright-fusion resonance solution, dark-fusion resonance solution, bright-fission resonance solution, and dark-fission resonance solution. By introducing two polynomials, the generalized rational solutions splitting from two peaks into three peaks are constructed. The dynamical behavior of these phenomena are demonstrated in detail. With the analysis of the first- to third-order rational solution, it is extremely interesting to find that s-order rational solution has 2s + 1 points for rogue wave-type solution or parallel lines for W-shaped solution. More importantly, the bright rational solution must have s peaks above the zero background plane and s + 1 valleys below the background plane, while the dark one is just the opposite.

The remainder of our article is organized as follows. In Section 2, bright and dark cases of rogue-type rational solution, W-type rational solution, high-order rational solutions and generalized rational solutions for the (3+1)-dimensional KS equation are constructed by virtue of the Hirota bilinear method. Section 3 is devoted to deriving four types of bright-fusion, dark-fusion, bright-fission, and dark-fission resonance solutions for the (3+1)-dimensional KS equation. Finally, some conclusions are given in Section 4.

2. Bright and dark high-order rational solutions

In this section, rogue wave-type rational solutions, W-shaped rational solutions, high-order rational solutions, and generalized rational solutions are investigated for the (3+1)-dimensional KS equation (4). Let $\xi = x + my + nt$ in Eq. (4). It follows that

$$\eta u_{\xi\xi} + \frac{1}{2} \alpha (u^2)_{\xi\xi} + \beta u_{\xi\xi\xi\xi} + \delta u_{zz} = 0, \qquad (5)$$

here $\eta = \gamma m^2 + n$. It is then deduced that

$$\begin{aligned} &(\eta D_{\xi}^{2} + \beta D_{\xi}^{4} + \delta D_{z}^{2})f \cdot f \\ &= \eta (ff_{\xi\xi} - f_{\xi}^{2}) + \beta (ff_{\xi\xi\xi\xi} - 4f_{\xi}f_{\xi\xi\xi}3f_{\xi\xi}^{2}) \\ &+ \delta (ff_{zz} - f_{z}^{2}) = 0, \end{aligned}$$
(6)

with the transformation

$$u = \frac{12\beta}{\alpha} (\ln f(\xi, z))_{\xi\xi}, \tag{7}$$

where D_{ξ} and D_z are Hirota operators.^[16]

In order to construct the rational solution (7) of the (3+1)dimensional KS equation (4), we first introduce a polynomial function $F_s = f$ with the following form:

$$F_{s}(\xi,z) = \sum_{j=0}^{s(s+1)/2} \sum_{i=0}^{j} a_{s(s+1)-2j,2i} z^{2i} \xi^{s(s+1)-2j}, \ s \in \mathbb{Z}^{+}.$$
 (8)

Substituting F_s into Eq. (6) and collecting the coefficients of ξ and z, we can get a set of equations that determine the value of $a_{i,j}$ by setting the coefficients equal to 0. According to the transformation (7), the specific rational solution of Eq. (4) is obtained. Obviously, the power of the denominator for the solution u with respect to the independent variables, is always higher than that of the numerator, which gives rise to $u \rightarrow 0$ as $x^2 + y^2 + z^2 + t^2 \rightarrow \infty$.

Remark 1 The proposed form of Eq. (8) has some limitations on constructing the nontrivial rational solutions of the nonlinear models in the bilinear form, which means it is not applicable to all the bilinear equations. Only all the bilinear operator in these models are linear combinations of D_v^{2p} ($p \in \mathbb{Z}^+$, v denotes independent variable) or they can be transformed into this form could be suitable for this method, otherwise the coefficients $a_{i,i}$ obtained are all trivial. If $a_{i,i}$ is trivial, then the first-order rational solution u = 0, let alone the high-order ones. Many examples can be calculated to verify the above conclusion, such as (2+1)-dimensional Benjamin–Ono equation.^[27] (2+1)-dimensional Camassa-Holm-Kadomtsev-Petviashvili equation,^[28] (3+1)-dimensional Hirota equation,^[29] (3+1)dimensional KP equation.^[30] and so on.

For s = 1, a set of non-trivial coefficients of function F_1 is obtained, *i.e.*, $a_{0,2} = -\eta^2 a_{0,0}/3\beta \delta$, $a_{2,0} = -\eta a_{0,0}/3\beta$. Plugging them into Eq. (7), there appears the expression for the first-order rational solution in the form

$$u_1(\xi, z) = -\frac{24\beta\delta\eta(3\beta\delta - 2\eta z^2 - \eta\delta\xi^2)}{\alpha(3\beta\delta - \eta^2 z^2 + \eta\delta\xi^2)^2},$$
(9)

where $\xi = x + my + nt$ and α , β , δ are all non-zero real numbers. Then we can get the following rogue wave-type and W-shaped rational solutions in case 1 and case 2.

Case 1 Rogue wave-type rational solutions

At the plane (y, z), we can obtain three critical points y = 0and $y = \pm (3/m)\sqrt{-\beta/\eta}$ along the line z = 0 through solving the following equations

$$\begin{cases} u_{1,y}(x=0,t=0) = 0\\ u_{1,z}(x=0,t=0) = 0. \end{cases}$$

Since $u_1(x = 0, t = 0, z = 0)$ is an even function about y, the values of u_1 are the same at the last two opposite points. At the point of (x = 0, t = 0, z = 0), the corresponding values of u_1 are $-8\eta/\alpha$ at the point y = 0 and η/α at the points $y = \pm (3/m) \sqrt{-\beta/\eta}$. After further analysis, it is clear that when the sigh of α , β are the same and η , δ are opposite, a bright rogue wave-type rational solution can be obtained. However, when the sigh of α , δ , η are the same, β is opposite of theirs, a dark rational solution is derived. It is not difficult to find that the dark rational solution also has three extreme points, which are exactly the same as that of bright rational solution. The corresponding amplitudes of u_1 are the same in the two situations, but the directions of the maximum amplitude point are opposite. As shown in Fig. 1, it illustrates the characteristics of the above rational solutions and the dynamic behavior. Red points denote the extreme values of the bright rational solution, while blue points represent that of the dark rational solution.



Fig. 1. The first-order rational solution of the (3+1)-dimensional KS equation (4) at the plane (y,z) with the parameters $(\beta, \gamma, \delta, m, n) = (1, -1, -1, 2, 2)$. (a) Bright rogue wave-type rational solution with $\alpha = 1$, (b) dark rogue wave-type rational solution with $\alpha = -1$.

Similarly, rogue wave-type rational solutions are also available for the planes (x,z) and (z,t). Along the line z = 0, there both exist three extreme points at the plane (x,z) for $u_1(y = 0, t = 0)$ and the plane (z,t) for $u_1(x = 0, y = 0)$,

which are at the points x = 0 and $x = \pm 3\sqrt{-\beta/\eta}$, t = 0 and $t = \pm (3/n)\sqrt{-\beta/\eta}$, respectively. The characteristic properties of these two planes are similar as the plane (y, z), here the dynamical behavior will not be elaborated. For the above three planes, whether bright or dark rational solutions, the corresponding values of u_1 are $-8\eta/\alpha$ and η/α at the extreme points, which are all proportional to η/α .

Case 2 W-shaped rational solutions

For the plane (x, y), three parallel lines can be obtained by the method of finding the extreme point as in case 1, which are x = -my, and $x = -my \pm 3\sqrt{-\beta/\eta}$, respectively. It provides us that

$$\begin{cases} u_1(x = -my, z = 0, t = 0) = -\frac{8\eta}{\alpha}, \\ u_1(x = -my \pm 3\sqrt{\frac{-\beta}{\eta}}, z = 0, t = 0) = \frac{\eta}{\alpha}. \end{cases}$$

The three parallel lines denote peaks or valleys of the Wshaped rational solution. According to the distance formula of two parallel lines, the width of the W-shaped rational solution is $6\sqrt{-\beta/\eta(1+m^2)}$. In terms of the expressions for the three parallel lines, it is not difficult to find that the absolute value of *m* determines the angle between the W-shaped rational solution and the positive half of the y axis, and its sigh determines the rotation direction. So as m changes, the W-shaped rational solution will have a corresponding tilt. The amplitude of u_1 is determined by γ , *m*, *n*, and α together. In addition, we can obtain different W-shaped solutions by changing the sign of the parameters, *i.e.*, α , β , and η . When the sigh of η is different from α and β , the W-shaped rational solution is a bright type. While the sigh of β is different from α and η , it yields a dark W-shaped rational solution. The dynamical phenomena of the bright and dark W-shaped rational solutions are demonstrated in Fig. 2.



Fig. 2. The first-order W-shaped rational solution of the (3+1)-dimensional KS equation (1) at (y,z) plane with the parameters $(\alpha,\beta,\gamma,\delta,m,n)$ are: (a) (1,1,-1,1,2,2); (b) (1,1,-1,1,-2,2); (c) (-1,1,-1,1,2,2); (d) (-1,1,-1,1,-2,2). (a) and (b) bright W-shaped rational solution; (c) and (d) dark W-shaped rational solution.

When y = z = 0, the solution u_1 has three parallel lines t = -x/n and $t = -x/n \pm (3/n)\sqrt{-\beta/\eta}$, their corresponding values are $-8\eta/\alpha$ and η/α , respectively. The parameter *n* controls the tilt of solution, and the width of last two parallel lines is $6\sqrt{-\beta/\eta(1+n^2)}$. At the (y,t) plane, there also have three parallel lines t = -m/ny and $t = -(m/n)y \pm (3/n)\sqrt{-\beta/\eta}$. The expression m/n also determines the tilt angle of solution, and the width of last two parallel lines is $6\sqrt{-\beta/\eta(m^2+n^2)}$.

From the analysis of the above two cases, the values of the first-order rational solutions are both $-8\eta/\alpha$ and η/α at the critical points or on the parallel lines, whether the rational solution is a W-shaped or rogue wave-type solution.

Case 3 High-order rational solutions

The above discussed are the cases of the first-order rational solutions, high-order rational solution can be derived with the increase of *s*. Here we take s = 2 and s = 3 as examples to analyze the dynamic characteristic of the high-order rational solution at (y,z) and (x,y) planes. For s = 2, a set of coefficients $a_{i,j}$ of the function F_2 can be derived, that is

$$\begin{cases} a_{0,0} = \frac{75\beta^2 a_{4,0}}{\eta^2}, \ a_{0,2} = -\frac{19\beta a_{4,0}}{\delta}, \ a_{0,4} = \frac{17\eta^2 a_{4,0}}{25\delta^2}, \\ a_{0,6} = -\frac{\eta^4 a_{4,0}}{25\beta\delta^3}, \ a_{2,0} = \frac{5\beta a_{4,0}}{\eta}, \ a_{2,2} = \frac{18\eta a_{4,0}}{5\delta}, \\ a_{2,4} = -\frac{3\eta^3 a_{4,0}}{25\beta\delta^2}, \ a_{4,2} = -\frac{3\eta^2 a_{4,0}}{25\beta\delta}, \ a_{6,0} = -\frac{\eta a_{4,0}}{25\beta} \end{cases}.$$

Taking the coefficients into Eq. (8), It then follows Eq. (7) that the expression of second-order rational solution $u_2(\xi, z)$ can be adduced, namely,

$$u_2(\xi, z) = \frac{P}{Q},\tag{10}$$

where

$$\begin{split} P &= -1440\beta^2\delta^5\eta^6\xi^6z^2 - 42000\beta^3\delta^6\eta^4\xi^6 + 144\delta^3\eta^9\beta\xi^4z^6 - 3384\beta^2\delta^2\eta^9z^8 + 67920\beta^3\delta^3\eta^7z^6 \\ &- 1110000\beta^4\delta^4\eta^5z^4 - 72\delta^6\eta^6\beta\xi^{10} - 216\delta^5\eta^7\beta\xi^8z^2 + 1800\beta^2\delta^6\eta^5\xi^8 + 2625000\beta^5\delta^5\eta^3z^2 \\ &+ 5625000\beta^6\delta^6\eta - 144\eta^8\beta\xi^6z^4 + 216\delta^2\eta^{10}\beta\xi^2z^8 + 2016\beta^2\delta^3\eta^8\xi^2z^6 + 90000\beta^3\delta^4\eta^6\xi^2z^4 \\ &- 3060000\beta^4\delta^5\eta^4\xi^2z^2 + 6375000\beta^5\delta^6\eta^2\xi^2 + 72\eta^{11}\delta\beta z^{10} + 2160\beta^2\delta^4\eta^7\xi^4z^4 \\ &+ 126000\beta^3\delta^5\eta^5\xi^4z^2 - 750000\beta^4\delta^6\eta^3\xi^4, \end{split}$$

$$\begin{aligned} Q &= 15\delta^2\eta^{10}\alpha\xi^4z^8 + 6\eta^{11}\delta\alpha\xi^2z^{10} + 468750\beta^5\delta^6\eta\alpha\xi^2 - 50\beta\delta^6\eta^5\alpha\xi^{10} + 15\delta^4\eta^8\alpha\xi^8z^4 + \delta^6\eta^6\alpha\xi^{12} \\ &+ 20\delta^3\eta^9\alpha\xi^6z^6 + 2500\beta^3\delta^6\eta^3\alpha\xi^6 + 109375\beta^4\delta^6\eta^2\alpha\xi^4 - 330\beta\delta^5\eta^6\alpha\xi^8z^2 - 724\beta\delta^4\eta^7\alpha\xi^6z^4 \\ &+ 4700\beta^2\delta^5\eta^5\alpha\xi^6z^2 - 282\beta\delta^2\eta^9\alpha\xi^2z^8 + 5660\beta^2\delta^3\eta^7\alpha\xi^2z^6 - 92500\beta^3\delta^4\eta^5\alpha\xi^2z^4 + \eta^{12}\alpha z^{12} \\ &- 692\beta\delta^3\eta^8\alpha\xi^4z^6 + 1239\beta^2\delta^2\eta^8\alpha z^8 - 12500\beta^3\delta^5\eta^4\alpha\xi^4z^2 + 3515625\beta^6\delta^6\alpha - 34\beta\delta\eta^{10}\alpha z^{10} \\ &+ 11050\beta^2\delta^4\eta^6\alpha\xi^4z^4 - 19900\beta^3\delta^3\eta^6\alpha z^6 + 289375\beta^4\delta^4\eta^4\alpha z^4 - 1781250\beta^5\delta^5\eta^2\alpha z^2 \\ &+ 218750\beta^4\delta^5\eta^3\alpha\xi^2z^2 + 375\beta^2\delta^6\eta^4\alpha\xi^8 + 6\delta^5\eta^7\alpha\xi^{10}z^2 \end{aligned}$$

When the parameters selected in u_2 are the same as those in u_1 of case 1, it will give rise to the second-order rogue wavetype rational solution, namely, bright and dark rational solution, as shown in Fig. 3. According to the analysis process of the first-order rational solution, we can also calculate the coordinates of the red and blue points in the (y, z) plane and their corresponding values of the second-order rational solutions. The y and z coordinates of the red points are approximately localized at $(\pm 1.768, 0)$, (0, 0), and $(\pm 0.821, 0)$. Their corresponding values of u_2 are approximately -3.2 for the first three points and 19.385 for the last two points. After further calculation, it is not difficult to find that the blue points and the red points have the same coordinates, while the corresponding values of u_2 are just opposite with each other. Figure 3(a) displays the dynamics of the second-order bright rogue wave-type rational solutions, while figure 3(b) shows the dark one.



Fig. 3. The second-order rogue wave-type rational solution in Eq. (4) at the (y,z) plane with the similar parameters as those given in Fig. 1. (a) Bright rogue wave-type rational solution; (b) dark rogue wave-type rational solution.

Similarly, when the parameters selected in u_2 are the same as those in u_1 of case 2, the second-order bright and dark W-shaped rational solutions can be constructed, as shown in Fig. 4. It is obvious that the tilt angles of the corresponding W-shaped rational solutions in Figs. 2 and 4 are the same and the slopes are $\pm 1/2$, which are determined by the parameter *m*.



Fig. 4. The second-order W-shaped rational solution in Eq. (4) at the (x,y) plane with the similar parameters as those given in Fig. 2. (a) Bright W-shaped rational solution; (b) dark W-shaped rational solution.

Setting s = 3 in Eq. (8), the expression for the coefficients $a_{i,j}$ in the function F_3 is as follows:

$$\begin{cases} a_{0,0} = -\frac{847\beta^2 a_{4,0}}{15\eta^2}, \ a_{0,2} = \frac{58\beta a_{4,0}}{\delta}, \\ a_{0,4} = -\frac{13381\eta^2 a_{4,0}}{4235\delta^2}, \ a_{0,6} = \frac{22828\eta^4 a_{4,0}}{148225\beta\delta^3}, \\ a_{0,8} = -\frac{2601\eta^6 a_{4,0}}{1037575\beta^2\delta^4}, \ a_{0,10} = \frac{174\eta^8 a_{4,0}}{5187875\beta^3\delta^5} \\ a_{0,12} = -\frac{3\eta^{10} a_{4,0}}{5187875\beta^4\delta^6}, \ a_{2,0} = \frac{154\beta a_{4,0}}{5\eta}, \\ a_{2,2} = -\frac{18\eta a_{4,0}}{55\delta}, \ a_{2,4} = -\frac{36\eta^3 a_{4,0}}{4235\beta\delta^2}, \\ a_{2,6} = -\frac{276\eta^5 a_{4,0}}{13475\beta^2\delta^3}, \ a_{2,8} = \frac{342\eta^7 a_{4,0}}{1037575\beta^3\delta^4}, \\ a_{2,10} = -\frac{18\eta^9 a_{4,0}}{5187875\beta^4\delta^5}, \ a_{4,2} = \frac{108\eta^2 a_{4,0}}{847\beta\delta}, \\ a_{4,4} = -\frac{642\eta^4 a_{4,0}}{29645\beta^2\delta^2}, \ a_{4,6} = \frac{876\eta^6 a_{4,0}}{1037575\beta^3\delta^3}, \\ a_{4,8} = -\frac{9\eta^8 a_{4,0}}{1037575\beta^4\delta^4}, \ a_{6,0} = \frac{4\eta a_{4,0}}{275\beta}, \\ a_{6,2} = -\frac{228\eta^3 a_{4,0}}{21175\beta^2\delta}, \ a_{6,4} = \frac{12\eta^5 a_{4,0}}{13475\beta^3\delta^2}, \\ a_{6,6} = -\frac{12\eta^7 a_{4,0}}{1037575\beta^4\delta^3}, \ a_{8,4} = -\frac{9\eta^6 a_{4,0}}{1037575\beta^4\delta^2}, \\ a_{10,0} = \frac{6\eta^3 a_{4,0}}{105875\beta^3}, \ a_{10,2} = -\frac{18\eta^5 a_{4,0}}{5187875\beta^4\delta}, \\ a_{12,0} = -\frac{3\eta^4 a_{4,0}}{5187875\beta^4} \right\}.$$

By taking them into Eq. (7) with Eq. (8), it yields the ex-

pression of the third-order rational solution $u_3(\xi, z)$. Since the expression of $u_3(\xi, z)$ is too cumbersome, we only shows its dynamic behavior, as illustrated in Figs. 5 and 6.



Fig. 5. The third-order rational solution at the (y, z) plane with the similar parameters as those given in Fig. 1. (a) Bright rogue wave-type rational solution; (b) dark rogue wave-type rational solution.



Fig. 6. The third-order W-shaped rational solution in Eq. (4) at the (x, y) plane with similar parameters as those given in Fig. 2. (a) and (b) Bright W-shaped rational solution; (c) and (d) dark W-shaped rational solution.

There are seven critical points for the third-order rational solution on the z = 0 axis at the (y,z) plane, and they also have the above symmetry results, see Fig. 5. The y and z coordinates of the critical points for the bright and dark thirdorder rogue wave-type rational solution are approximately localized at $(\pm 2.54, 0)$, $(\pm 1.565, 0)$, $(\pm 0.718, 0)$, (0, 0), and the corresponding values of u_3 are ± 3.534 , ± 18.556 , ± 4.516 , ± 26.182 , respectively. Simultaneously, the third-order Wshaped rational solution has the same characteristics as the lower one, and seven critical lines can be seen in Fig. 6.

As *s* increases, high-order rational solution (rogue wavetype or W-shaped) of the (3+1)-dimensional KS equation can be generated. It is noticeable that the solution meets a rule: *s*-order rational solution has 2s + 1 points or parallel lines on different planes, and the two situations show different dynamic behavior characteristics.

Case 4 Generalized rational solutions

Attention is now turned to investigate generalized rational solutions for the (3+1)-dimensional KS equation. To achieve

this purpose, we introduce two polynomials

$$G_{s}(\xi,z) = \sum_{j=0}^{s(s+1)/2} \sum_{i=0}^{j} b_{s(s+1)-2j,2i} \xi^{2i} z^{s(s+1)-2j},$$

$$H_{s}(\xi,z) = \sum_{j=0}^{s(s+1)/2} \sum_{i=0}^{j} c_{s(s+1)-2j,2i} z^{2i} \xi^{s(s+1)-2j}, \quad (11)$$

in the same form as $F_s(\xi, z)$ in Eq. (8), where the two coefficients $b_{s(s+1)-2j,2i}$ and $c_{s(s+1)-2j,2i}$ will be determined later. Applied the method used in Ref. [26], it is natural to have the following result.

Theorem 1 The (3+1)-dimensional KS equation (4) has generalized rational solutions in the form

$$u_s(\xi, z; M, N) = \frac{12\beta}{\alpha} (\ln f(\xi, z; M, N))_{\xi\xi}, \qquad (12)$$

for $s \ge 1$, with

$$f = L_s(\xi, z; M, N)$$

= $F_{s+1}(\xi, z) + 2MzG_s(\xi, z) + 2N\xi H_s(\xi, z)$
+ $(M^2 + N^2)F_{s-1}(\xi, z),$ (13)

where M and N are arbitrary constants.

Obviously, let M, N, and s in Eq. (13) be equal to zero, the one-order rational solution can be obtained, which is consistent with the results obtained at the beginning of this section. Without loss of generality, when s = 1, a set of non-trivial co-

efficients of function L_1 can be obtained as

$$\begin{cases} a_{0,0} = -\frac{25\beta (M^2 \delta b_{0,2}^2 + 9N^2 \eta c_{2,0}^2 - 27\beta a_{4,0}^2)}{9(M^2 + N^2 + 1)\eta^2 a_{4,0}}, \\ a_{0,2} = -\frac{19\beta a_{4,0}}{\delta}, \ a_{0,4} = \frac{17\eta^2 a_{4,0}}{25\delta^2}, \\ a_{0,6} = -\frac{\eta^4 a_{4,0}}{25\beta\delta^3}, \ a_{2,0} = \frac{5\beta a_{4,0}}{\eta}, \ a_{2,2} = \frac{18\eta a_{4,0}}{5\delta}, \\ a_{2,4} = -\frac{3\eta^3 a_{4,0}}{25\delta^2\beta}, \ a_{4,2} = -\frac{3\eta^2 a_{4,0}}{25\beta\delta}, \\ a_{6,0} = -\frac{\eta a_{4,0}}{25\beta}, \ b_{0,0} = -\frac{5\beta b_{0,2}}{3\eta}, \ b_{2,0} = -\frac{\eta b_{0,2}}{3\delta}, \\ c_{0,0} = \frac{\beta c_{2,0}}{\eta}, \ c_{0,2} = -\frac{3\eta c_{2,0}}{\delta} \\ \end{cases}.$$

Plugging them into Eq. (7), it gives rise to the first generalized rational solution $u_2(\xi, z; M, N)$. Here we only analyze their dynamic behavior.

When M = N = 0, it becomes the obtained rational solution and can bring in similar results as shown in Fig. 3 by selecting appropriate values for the remaining parameters. As |M| and |N| increase, the solution can be split from two peak into three peaks rogue wave solution, which is also known as 'rogue wave triplet' solution. Figure 7 shows its evolution process of the bright case. Let $\alpha = -1$, the dark case can be obtained. Using Theorem 1, high-order generalized rational solutions can be derived for the case of $s \ge 2$.



Fig. 7. Decomposition process of the generalized rational solution $u_2(\xi, z; M, N)$ for the (3+1)-dimensional KS equation with distinct values of the parameters M, N and $(\alpha, \beta, \gamma, \delta, m, n, a_{4,0}, b_{0,2}, c_{2,0}) = (1, 1, -1, -1, 2, 2, 2, 2, 2)$. (a): M = N = 0; (b): M = N = 1; (c): M = N = 3; (d): M = N = 5; (e): M = N = 8; (f): M = N = 10.

3. The *N*-wave resonance solution

$$(D_x D_t + \beta D_x^4 + \gamma D_y^2 + \delta D_z^2) f \cdot f = 0, \qquad (14)$$

The purpose of this section is to construct *N*-wave resonance solutions of the (3+1)-dimensional KS equation (4) by reverting it into the following bilinear form:

where D is the Hirota operator,^[16] here use has been made of the transformation

$$u = \frac{12\beta}{\alpha} (\ln f(x, y, z, t))_{xx}.$$
 (15)

Assuming

$$f = 1 + \sum_{i=1}^{N} e^{k_i x + l_i y + m_i z + w_i t}$$

and then plugging it into Eq. (14), one can have the dispersion relation

$$w_i = -\frac{\beta k_i^4 + \gamma l_i^2 + \delta m_i^2}{k_i}.$$
(16)

Let $l_i = ak_i^2$, $m_i = bk_i^2$. It thus transpires that *f* has the following specific expression

$$f = 1 + \sum_{i=1}^{N} e^{k_i x + ak_i^2 y + bk_i^2 z - 4\beta k_i^3 t},$$
(17)

with an additional constrain condition $3\beta = \gamma a^2 + \delta b^2$, which gives the solution of Eq. (14). Substituting Eq. (17) into Eq. (15), the *N*-wave resonance soliton solution can be derived. For the soliton solution of Eq. (4), it exists both fission and fusion phenomena.

By choosing appropriate parameters, four different resonance solutions are discussed, including bright-fission, darkfission, bright-fusion, and dark-fusion solutions. Figure 8 displays different dynamical phenomena of the two-wave resonance solution. Fission waves can be split from one wave to multiple waves, while fusion waves are just the opposite. In order to understand the fusion and fission processes of the soliton solutions clearly, the three-wave resonance solution is taken as an example, and the evolution processes of fusion and fission over time are given respectively. Figure 9 shows the bright type of the three-wave resonance solution, dark type of the three-wave resonance solution also has the same phenomena. In the same way, *N*-wave resonance solution can be constructed.



Fig. 8. Two-wave resonance solution of the (3+1)-dimensional KS equation (4) in the (*x*,*z*) plane with parameters (α , β , *a*, *b*, γ , δ , *k*_1, *k*_2) as: (a) (1, 1, 1, 1, $\sqrt{2}$, 1, 1, -1/2, -1) bright-fusion solution; (b) (-1, 1, 1, 1, $\sqrt{2}$, 1, 1, -1/2, -1) dark-fusion solution; (c) (1, 1, 1, 1, $-\sqrt{2}$, 1, 1, 1/2, 1) bright-fission solution; (d) (-1, 1, 1, 1, $-\sqrt{2}$, 1, 1, 1/2, 1) dark-fission solution.



Fig. 9. The evolution of three-wave resonance solution over time with similar parameters as those given Figs. 8(a) and 8(c). Top row: bright-fusion solution ($k_3 = -3/2$). (d)–(f) bright-fission solution ($k_3 = 3/2$).

4. Summary and discussion

A (3+1)-dimensional Kudryashov–Sinelshchikov equation modelling the liquid containing gas bubbles has been studied in this paper. Rogue wave-type rational solution, W-shaped rational solution, Multiple basic rogue wave solution, and *N*wave resonance solution are provided explicitly for this equation based on two kinds of bilinear equations.

By introducing the polynomial function (8), we obtain bright rogue wave-type rational solution, dark rogue wavetype rational solution, bright W-shaped rational solution, and dark W-shaped rational solution, see Figs. 1–6. From the analysis of the first- to third-order rational solutions, it is not difficult to find that *s*-order rational solution has 2s + 1 points for rogue wave-type solution or parallel lines for W-shaped solution. As long as it is a bright rational solution, there must be *s* peaks above the background plane (zero plane), s + 1 valleys below the background plane, and the dark one is just the opposite. Their corresponding properties, such as amplitude, wave width, and tilt angle, are all parametrically controllable. Moreover, generalized rational solutions splitting from two peaks to three peaks are derived by introducing two polynomials (11). The dynamical behavior of the splitting process has been demonstrated in Fig. 7.

In addition, the *N*-wave resonance solution of the (3+1)dimensional KS equation is constructed under the dispersion relation and specific constraints. Bright-fission, dark-fission, bright-fusion, and dark-fusion two-wave resonance solutions are constructed, see Fig. 8. Fission waves can be split from one wave to multiple waves, while fusion waves are just the opposite. Figure 9 shows the evolution processes of fusion and fission three-wave resonance solutions, which reveal diverse dynamic behavior.

References

- [1] Kudryashov N A and Sinelshchikov D I 2012 Phys. Scr. 85 025402
- [2] Ma W X 1993 J. Phys. A: Math. Gen. 26 L17
- [3] Lax P D 1968 Commun. Pure Appl. Math. 21 467

- [4] Kadomtsev B B and Petviashvili V I 1970 Sov. Phys. Dokl. 15 539
- [5] Kudryashov N A and Sinelshchikov D I 2010 Phys. Lett. A 374 2011
- [6] Bona J L and Schonbek M E 1985 Proc. R. Soc. Edinburgh 101A 207
- [7] Wazwaz A M 2008 Appl. Math. Comput. 200 437
- [8] Mohammed K 2009 Commun. Nonlinear Sci. Numer. Simulat. 14 1169
- [9] Yue Y F, Huang L L and Chen Y 2020 Commun. Nonlinear Sci. Numer. Simulat. 89 105284
- [10] Yan Z Y 2017 Chaos 27 053117
- [11] Wang Y H and Wang H 2017 Nonlinear Dyn. 89 235
- [12] Wang Y H and Wang H 2014 Phys. Scr. 89 125203
- [13] Yue Y F, Huang L L and Chen Y 2019 Appl. Math. Lett. 89 70
- [14] Jin X W and Lin J 2020 J. Magn. Magn. Mater. 502 166590
- [15] Chen Y, Huang L L and Liu Y 2020 J. Nonlinear Sci. 30 93
- [16] Hirota R 2004 The direct method in soliton theory (Cambridge: Cambridge University Press)
- [17] Guo B L, Ling L M and Liu Q P 2012 Phys. Rev. E 85 026607
- [18] Chen J C, Chen L Y, Feng B F and Maruno K I 2019 Phys. Rev. E 100 052216
- [19] Huang L L and Chen Y 2018 Nonlinear Dyn. 92 221
- [20] Zhou A J and Chen A H 2018 Phys. Scripta 93 125201
- [21] Feng Y L, Shan W R, Sun W R, et al. 2014 Commun. Nonlinear Sci. Numer. Simulat. 19 880
- [22] Tu J M, Tian S F, Xu M J, et al. 2016 Nonlinear Dyn. 83 1199
- [23] Liu W H and Zhang Y F 2018 Mod. Phys. Lett. B 32 1850359
- [24] Liu W H and Zhang Y F 2020 Waves Random Complex Media 30 470
- [25] Wang M, Tian B, Sun Y, et al. 2020 Comput. Math. Appl. 79 576
- [26] Clarkson P A and Dowie E 2017 Transactions of Mathematics and its Applications 1 1 tnx003
- [27] Zhao Z L, He L C and Gao Y B 2019 Complexity bf 2019 8249635
- [28] Qin C Y, Tian S F, Wang X B, et al. 2018 Commun. Nonlinear Sci. Numer. Simulat. 62 378
- [29] Liu W H and Zhang Y F 2019 Appl. Math. Lett. 98 184
- [30] Cui W Y and Zha Q L 2018 Comput. Math. Appl. 76 1099