

A LETTERS JOURNAL EXPLORING THE FRONTIERS OF PHYSICS

OFFPRINT

Explosive oscillation death in coupled Stuart-Landau oscillators

Hongjie Bi, Xin Hu, Xiyun Zhang, Yong Zou, Zonghua Liu and Shuguang Guan

EPL, 108 (2014) 50003

Please visit the website
www.epljournal.org

Note that the author(s) has the following rights:

- immediately after publication, to use all or part of the article without revision or modification, **including the EPLA-formatted version**, for personal compilations and use only;

- no sooner than 12 months from the date of first publication, to include the accepted manuscript (all or part), **but not the EPLA-formatted version**, on institute repositories or third-party websites provided a link to the online EPL abstract or EPL homepage is included.

For complete copyright details see: https://authors.epletters.net/documents/copyright.pdf.

THE FRONTIERS OF PHYSICS AN INVITATION TO SUBMIT YOUR WORK

A LETTERS JOURNAL EXPLORING

www.epljournal.org

The Editorial Board invites you to submit your letters to EPL

EPL is a leading international journal publishing original, innovative Letters in all areas of physics, ranging from condensed matter topics and interdisciplinary research to astrophysics, geophysics, plasma and fusion sciences, including those with application potential.

The high profile of the journal combined with the excellent scientific quality of the articles ensures that EPL is an essential resource for its worldwide audience. EPL offers authors global visibility and a great opportunity to share their work with others across the whole of the physics community.

Run by active scientists, for scientists

EPL is reviewed by scientists for scientists, to serve and support the international scientific community. The Editorial Board is a team of active research scientists with an expert understanding of the needs of both authors and researchers.



www.epljournal.org

A Letters Journal Exploring the Frontiers of Physics







10,755 citations in 2013

"We greatly appreciate the efficient, professional and rapid processing of our paper by your team."

Cong Lin Shanghai University

Six good reasons to publish with EPL

We want to work with you to gain recognition for your research through worldwide visibility and high citations. As an EPL author, you will benefit from:



Quality – The 50+ Co-editors, who are experts in their field, oversee the entire peer-review process, from selection of the referees to making all final acceptance decisions.



Convenience – Easy to access compilations of recent articles in specific narrow fields available on the website.



Speed of processing – We aim to provide you with a quick and efficient service; the median time from submission to online publication is under 100 days.



High visibility – Strong promotion and visibility through material available at over 300 events annually, distributed via e-mail, and targeted mailshot newsletters.



International reach – Over 2600 institutions have access to EPL, enabling your work to be read by your peers in 90 countries.



Open access – Articles are offered open access for a one-off author payment; green open access on all others with a 12-month embargo.

Details on preparing, submitting and tracking the progress of your manuscript from submission to acceptance are available on the EPL submission website **www.epletters.net**.

If you would like further information about our author service or EPL in general, please visit **www.epljournal.org** or e-mail us at **info@epljournal.org**.

EPL is published in partnership with:









IOP Publishing

www.epljournal.org



www.epljournal.org

Explosive oscillation death in coupled Stuart-Landau oscillators

HONGJIE BI¹, XIN HU¹, XIYUN ZHANG¹, YONG ZOU^{1,2}, ZONGHUA LIU^{1,2} and SHUGUANG GUAN^{1,2}

¹ Department of Physics, East China Normal University - Shanghai, 200241, China

² State Key Laboratory of Theoretical Physics, Institute of Theoretical Physics, Chinese Academy of Sciences Beijing 100190, China

received 21 August 2014; accepted in final form 7 November 2014 published online 1 December 2014

PACS 05.45.Xt - Synchronization; coupled oscillators PACS 89.75.Kd - Patterns

Abstract – Recently, explosive phase transitions, such as explosive percolation and explosive synchronization, have attracted extensive research interest. So far, most existing works have investigated Kuramoto-type models, where only phase variables are involved. Here, we report the occurrence of explosive oscillation quenching in a system of coupled Stuart-Landau oscillators that incorporates both phase and amplitude dynamics. We observe three typical scenarios with distinct microscopic mechanism of occurrence, *i.e.*, ordinary, hierarchical, and cluster explosive oscillation death, corresponding to different frequency distributions of oscillators. We carry out theoretical analyses and obtain the backward transition point, which is shown to be independent of the specific frequency distributions. Numerical results are consistent with the theoretical predictions.

Copyright © EPLA, 2014

Introduction. – Recently, two remarkable phenomena, *i.e.*, explosive percolation (EP) and explosive synchronization (ES), have attracted great attention in the area of nonlinear dynamics and complex networks.

EP refers to the sudden formation of an inter-connected giant cluster in a growing network, whose size is comparable with that of the whole network [1–9]. It is shown that EP could occur in a network under Achlioptas-like growth rules, which generally enhance the formation of small clusters and suppress the formation of giant clusters before transition [1]. Such processes are essentially different from the traditional percolation transition in the Erdős-Rényi network, where the size of the giant cluster grows gradually and continuously after the transition.

ES is an abrupt, irreversible synchronization first reported in a generalized Kuramoto model on a scale-free network when the natural frequency and degree of the *j*-th oscillator in the network are correlated, *i.e.*, $\omega_j = k_j$ [10]. This result is drastically different from the continuous synchronization transition found in the Kuramoto model for decades, and thus has stimulated intensive research along this line [11–18].

So far, ES is only observed in phase synchronization, either in the Kuramoto-type model or in the coupled Rössler system. One important question is: could explosive behaviors occur for the amplitude dynamics in networked oscillators? Motivated by this idea, we investigate a model of coupled Stuart-Landau (SL) oscillators in this paper. By introducing a similar coupling scheme as in ref. [16], we found that explosive oscillation death (EOD) could occur in the model. Specifically, we have numerically identified three typical types of EOD, namely, ordinary EOD (OEOD), hierarchical EOD (HEOD), and cluster EOD (CEOD), corresponding to different distributions of the natural frequency of SL oscillators. Interestingly, for all three cases, it is found that the backward phase transitions are independent of the specific frequency distributions (FDs). We further apply a self-consistency analysis, which proves the universal property of the backward phase transitions in the model.

The dynamical model. – Amplitude death (AD) and oscillation death (OD) both refer to the complete suppression of oscillations in coupled systems, *i.e.*, converting from oscillating states, such as limit cycles or chaos, into fixed points [19,20]. In AD all oscillators finally settle on the same fixed point, while in OD they settle on different fixed points. Since its first discovery in an array of coupled oscillators [21], AD and OD have stimulated continuous research in both theory [22–26] and experiments [27,28] in the past two decades. For networked oscillators, it is important to investigate how transition to AD or OD occurs, *i.e.*, does it occur suddenly for all oscillators, or does it develop gradually in the system? This issue has been addressed by several previous works. In refs. [29–31], it

FD	Formula	Parameter	K_f	K_b
Triangle	$g(\omega) = (\pi \Delta - \omega)/(\pi \Delta)^2,$	$\Delta = 0.1$	$K_f \approx 2.30$	$K_b \approx 2.02$
	$ \omega < \pi \Delta, 0$ otherwise			
Lorentzian	$g(\omega) = \frac{\Delta}{\pi(\omega^2 + \Delta^2)}$	$\Delta = 0.02$	$K_f \approx 3.32$	$K_b \approx 2.02$
Uniform	$g(\omega) = 1/(2\pi\Delta),$	$\Delta = 0.1$	$K_f^1 pprox 1.64$	$K_b^1\approx 2.02$
	$ \omega < \pi \Delta, 0$ otherwise		$K_f^2\approx 2.02$	$K_b^2 \approx 1.56$

Table 1: Summary of the three typical FDs with parameters investigated in this paper, and the critical points for both forward (K_f) and backward (K_b) transitions.

is found that with the increase of the coupling strength, partial oscillators in the network first undergo AD. Then more and more oscillators convert into the AD state as the coupling strength further increases until all oscillators finally become fixed points. In a few words, AD/OD occurs successively in such networked systems. Moreover, in ref. [32], although AD is shown to occur suddenly for all oscillators in the network, no hysteresis has been reported there. In this work, our primary concern is whether AD/OD could occur in an explosive way accompanied by a hysteresis in a certain coupled oscillators system. To this end, we study a dynamical model of networked SL oscillators with frequency-weighted coupling [16], *i.e.*,

$$\dot{z}_j(t) = (a + iw_j - |z_j|^2)z_j(t) + \frac{K|\omega_j|}{N} \sum_{n=1}^N [z_n(t) - z_j(t)].$$
(1)

Here $j = 1, 2, \dots, N$ is the index of oscillators. $z_j(t) = x_j(t) + iy_j(t)$ is the complex amplitude of the *j*-th oscillator at time *t*, and the dot represents the time derivative. *a* is a control parameter for the individual SL oscillator, *i.e.*, the dynamics settles on a limit cycle if a > 0, and on a fixed point if a < 0. ω_j is the natural frequency of the *j*-th oscillator, and *K* the uniform coupling strength. The most important characteristic of this model is that the effective coupling for oscillator *j* is proportional to its natural frequency ω_j . Therefore, the effective couplings in eq. (1) are heterogeneous rather than homogeneous as in most previous models.

In this work, we consider several typical FDs as listed in table 1. Without losing generality, we set a = 1 and only consider the situation of global coupling. Throughout this paper, numerical integration is carried out by the fourthorder Runge-Kutta method with time step 0.01. The initial phases of the limit cycles are random, *i.e.*, oscillators are uniformly distributed on the unit circle in a complex plane at the beginning.

The numerical results. – To characterize the collective behaviors of the coupled SL oscillators, two order parameters can be defined as: $Re^{i\psi} = \sum_{j=1}^{N} z_j(t)/N$, and $R_{\theta}e^{i\phi} = \sum_{j=1}^{N} e^{i\theta_j}/N$. Here, θ_j represents the phase of the *j*-th oscillator. The order parameter R ($0 \le R \le 1$ due to a = 1 in this study) characterizes the coherence



Fig. 1: (Color online) Characterization of the EOD by order parameters R and R_{θ} for three typical FDs, *i.e.*, triangle (left column), Lorentzian (middle column), and uniform (right column), respectively. System size N = 500. In both forward and backward transitions, the dynamical equations are integrated in an adiabatic way, where K is increased at a step of 0.02 and the final state for a prior K is used as the initial state for the next K. For each K, the order parameters are averaged in a time window after the transient stage. Such numerical schemes are adopted throughout this paper.

of the complete dynamics, including both amplitude and phase. The order parameter R_{θ} ($0 \le R_{\theta} \le 1$) only characterizes the phase coherence of the system, which does not involve any information of amplitude.

We now report the main numerical findings. Generally, we observe EOD in eq. (1), which, to our knowledge, is the first explosive phase transition involving the behavior of amplitude in a system of coupled oscillators. As listed in table 1, we consider three typical FDs in eq. (1). In fig. 1, we plot the order parameters R and R_{θ} vs. the coupling strength K. For both R and R_{θ} , we observe a sudden, discontinuous jump when the coupling strength K exceeds the critical value, *i.e.*, the forward transition point. After the transition, the order parameters are significantly greater than 0, indicating that the system goes from the incoherent state into the (partially) coherent one. Inversely, when the system starts from a coherent state initially, as the coupling strength decreases, the order parameters also experience an abrupt fall, but at a different coupling



Fig. 2: (Color online) Classification of the EOD by three measures: the effective frequencies Ω_j ; N_d/N (N_c/N); and the relative phases η_j (see the definitions in the text). For a better visualization, we have applied downsampling in this figure when necessary. All parameters are the same as in fig. 1. In our study, we have also carefully examined the process for the backward transition (not shown). Except for the transition points, the physical pictures are qualitatively the same.

strength from the forward transition point. In this way, a hysteresis loop, which is a typical characteristic in the first-order phase transitions, can be defined by the backward and forward transition points. It should be pointed out that, at this stage, we cannot identify whether it is OD or AD only based on the results in fig. 1. Later, we will provide more evidences to confirm that it is OD rather than AD that occurs in this system.

Interestingly, we further reveal that the phenomena of EOD shown in fig. 1 actually correspond to three qualitatively different situations, namely, the OEOD, the HEOD, and the CEOD. To characterize them, we define several quantitative measures. The first is the effective frequency for each oscillator: $\Omega_j = \frac{1}{T} \int_t^{t+T} \dot{\theta}_j(\tau) d\tau$ with $T \gg 1$, which basically is the averaged instantaneous frequency. The second is the relative phase with respect to the average phase: $\eta_j = \theta_j - \phi$, where ϕ is defined as in the order parameter R_{θ} . The third are two ratios N_d/N and N_c/N , where N_d and N_c are the numbers of oscillators that are in the OD state and the clustering state, respectively. In fig. 2, we plot these measures for the forward transition processes. In the following, we describe these three types of EOD in detail.

1) The OEOD. This corresponds to the triangle FD, and is basically the same as the situation of ES observed in previous works [10–12,16]. As shown in figs. 2(a1) and 2(a2), with the increase of K, oscillators oscillate almost according to their natural frequencies. This situation continues until K arrives at the forward transition point, where all the effective frequencies of oscillators suddenly and simultaneously collapse to 0. Note that 0 is the averaged natural frequency due to the symmetry of FDs in this study. This transition point characterizes the occurrence of EOD in the system. Moreover, as shown in fig. 2(a3), after the OD, the oscillators settle on different fixed points in the phase space that form two clusters.

- 2) The HEOD. This corresponds to the Lorentzian FD. As shown in figs. 2(b1) and 2(b2), with the increase of K, the effective frequencies of some oscillators collapse hierarchically to small values that are very close to 0. This process starts from the largest frequencies and monotonically moves to smaller frequencies. A careful examination of the dynamics of individual oscillator reveals that those oscillators whose effective frequencies have collapsed actually form two small clusters, rotating very slowly near the origin on the complex plane (fig. 3(b2)). However, since this collapse of effective frequencies before the forward transition point only involves minority oscillators in the system (fig. 2(b2)), their contributions to the order parameters can be ignored. Therefore, the order parameters have no significant growth at this stage, as shown in figs. 1(b1) and 1(b2). This situation remains until K arrives at the forward transition point, where the effective frequencies of all oscillators suddenly collapse to 0. Since all oscillators simultaneously become OD (fig. 2(b2)), the order parameter behaves in an explosive way, as shown in figs. 1(b1)and 1(b2).
- 3) The CEOD. This corresponds to the uniform FD. As shown in figs. 2(c1) and 2(c2), with the increase of K, all oscillators oscillate almost according to their natural frequencies before the first transition point $(K_f^1 \approx 1.64)$. Then when K arrives at this point, those oscillators with relatively large natural frequencies suddenly become locked and form two clusters. Inside each cluster, all oscillators have the same effective frequency, and they behave like two (giant) oscillators, rotating clockwise and counterclockwise, respectively. Since the number of oscillators involved in this clustering is comparable to the system size (fig. 2(c2)), the order parameters jump explosively, as shown in figs. 1(c1) and 1(c2). After the first transition, the system is in a partially coherent state, where clusters coexist with drifting oscillators. When K is increased further, the two clusters rotate slower and slower. Meanwhile, they gradually absorb the drifting oscillators until finally the effective frequencies of the two giant clusters completely approach 0 at the second critical point $(K_f^2 \approx 2.02)$, where OD occurs. So in this case, the system undergoes two transitions: one is explosive (at $K_f^1 \approx 1.64$); and the other is continuous (at $K_f^2 \approx 2.02$). The transition involves three distinct dynamical states, *i.e.*, the incoherent state,



Fig. 3: (Color online) Characterization of the EOD in the phase space, corresponding to three FDs, *i.e.*, triangle (left column), Lorentzian (middle column), and uniform (right column), respectively. In the first row, the amplitudes $|z_j|$, averaged for a long time after transient stage, are plotted vs. K. The other panels are the snapshots of system states corresponding to different K. The second and the third rows correspond to the states just before and after the first transitions, respectively. In (a2), (b2), (c2), and (c3), red/pinkish-red (blue/green) dots correspond to oscillators rotating clockwise (counterclockwise). In (a3), (b3), (a4)–(c4), red (blue) squares indicate the fixed points converted from oscillators with positive (negative) natural frequencies. The unit circle is plotted to guide the eyes. All parameters are the same as in fig. 1.

the partially coherent state, and the OD state. We emphasize that from figs. 1(c1) and 1(c2), the order parameters actually change continuously when OD occurs at the second critical point. However, since there is an explosive transition in the way toward OD in this case, we still term the whole transition process as CEOD.

In fig. 2, we have characterized the observed EOD from the perspective of phase, ignoring the amplitudes of oscillators. To completely illustrate the physical picture in the above three EOD processes, we now further characterize them in the phase space, as shown in fig. 3. Note that in our numerical simulations, initially all oscillators are randomly distributed on the unit circle in the complex plane.

In the case of OEOD, with the increase of K, oscillators gradually move inside the unit circle, as shown in figs. 3(a1) and 3(a2). However, according to fig. 2(a1),

these oscillators still rotate almost according to their natural frequencies though their amplitudes have been suppressed. Then when K arrives at the forward critical point, all oscillations suddenly cease and OD occurs, as shown in figs. 3(a2) and 3(a3). After OD, two compact clusters consisting of different fixed points are formed. Oscillators inside each cluster may slightly change positions when K further increases. However, oscillators cannot redistribute between two branches [33].

In the case of HEOD, from fig. 2(b1), we know that the effective frequencies of oscillators hierarchically collapse (near 0), starting from the largest frequency to the smaller ones successively. As shown in fig. 3(b2), we found that those oscillators form two small clusters near the origin. In the meantime, many oscillators move inside the unit circle, *i.e.*, their amplitudes are suppressed. Then when K exceeds the forward critical point, all oscillations in the system suddenly stop, indicating the occurrence of OD. Similarly, two compact clusters of fixed points form after OD.

In the case of CEOD, with the increase of K, oscillators gradually move inside the unit circle, *i.e.*, the oscillations are significantly suppressed. Even so, they still rotate almost according to their natural frequencies. Then when Kexceeds the first critical point $(K_f^1 \approx 1.64)$, some oscillators, *i.e.*, those with relatively large frequencies, suddenly become frequency-locked and form two clusters. After that, the system enters the partially coherent state, as shown in fig. 3(c3). Remarkably, it is clustering rather than OD that occurs at this point, which is essentially different from the two cases discussed above. When Kfurther increases, more and more drifting oscillators are recruited into the coherent clusters. Meanwhile, the rotations of the two clusters become slower and slower. Finally, when K exceeds the second critical point $(K_f^2 \approx 2.02)$, OD occurs, as shown in fig. 3(c4).

We emphasize that, because oscillators go to different fixed points in the quenched state, as shown in the inset of fig. 1(a1), it is OD rather than AD that occurs in this system. Moreover, in all three cases, with the further increase of K after OD, the formed two clusters will become more compact and they will more approach the unit circle, as shown in fig. 3.

The theoretical analysis. – In the previous section, three typical cases of EOD in networked SL oscillators have been characterized. In all three cases, the hysteresis loops are observed in phase diagrams. Particularly, it is found that a backward transition always occurs at $K_b \approx 2.02$ (fig. 1 and table 1), despite different FDs are used. Theoretically, it is desirable to predict both the forward and backward transition points for the EOD. However, the mathematical treatment turns out to be difficult even for the Kuramoto model, which only involves phase variables [12,17]. In the following we apply a selfconsistent analysis to obtain the backward transition point and prove that it is independent of specific FDs. To facilitate the analysis, we consider eq. (1) in polar coordinates, *i.e.*,

$$\dot{r}_j = (1 - r_j^2 - K|\omega_j|)r_j + \frac{K|\omega_j|}{N} \sum_{n=1}^N r_n \cos(\theta_n - \theta_j),$$

$$\dot{\theta}_j = \omega_j + \frac{K|\omega_j|}{N} \sum_{n=1}^N \frac{r_n}{r_j} \sin(\theta_n - \theta_j).$$
 (2)

Using the definition of order parameter, the above equation can be written in the mean-field form:

$$\dot{r}_j = (1 - r_j^2 - K|\omega_j|)r_j + KR|\omega_j|\cos(\psi - \theta_j),$$

$$\dot{\theta}_j = \omega_j + \frac{KR|\omega_j|}{r_j}\sin(\psi - \theta_j).$$
 (3)

We set a reference rotating frame, *i.e.*, $\psi(t) = \psi(0) + \langle \Omega \rangle t$, where $\langle \Omega \rangle$ is the average effective frequency of oscillators in the system. For a symmetric distribution of $g(\omega)$, we generally have $\langle \Omega \rangle = 0$. Defining $\eta_j \equiv \theta_j - \psi$, eqs. (3) become

$$\dot{r}_j = (1 - r_j^2 - K|\omega_j|)r_j + KR|\omega_j|\cos\eta_j,$$

$$\dot{\eta}_j = \omega_j - \frac{KR|\omega_j|}{r_j}\sin\eta_j.$$
 (4)

When all oscillators are phase-locked, we have $\dot{\eta}_j = 0$, then the stationary state for η_j can be solved as

$$\eta_j^* = \begin{cases} \eta_+^* = \arcsin(\frac{r_j}{KR}), & \omega_j > 0, \\ \eta_-^* = -\arcsin(\frac{r_j}{KR}), & \omega_j < 0. \end{cases}$$
(5)

Equation (5) implies that, in the phase-locked state, the oscillators in the system will evenly split into two clusters which are symmetric with respect to the real axis in the complex plane. If the coupling strength K further increases, they both approach the real axis gradually. Only in the limit case of $K \to \infty$, the two clusters coincide into one. This result has been numerically verified as shown in fig. 3.

Noticing that the system will form two very compact clusters in the phase space when K is large enough, we can approximately regard the whole system as two oscillators at the centroids of these two clusters. They are located at (r, η_+) and (r, η_-) , rotating with frequency $\omega > 0$ and $-\omega < 0$, respectively. Then eqs. (2) become

$$\dot{r} = (1 - r^2 - K\omega)r + K\omega r [1 + \cos(\eta_+ - \eta_-)]/2, \dot{\eta}_+ = \omega + K\omega \sin(\eta_- - \eta_+)/2, \dot{\eta}_- = -\omega + K\omega \sin(\eta_+ - \eta_-)/2.$$
(6)

By setting $\Theta = \eta_+ - \eta_-$, we obtain

$$\dot{r} = (1 - r^2 - K\omega)r + K\omega r[1 + \cos\Theta]/2,$$

$$\dot{\Theta} = 2\omega - K\omega \sin\Theta.$$
(7)

From the above equations, Θ has steady solution Θ^* only when $K \ge 2$, *i.e.*,

$$\sin \Theta^* = 2/K,\tag{8}$$



Fig. 4: (Color online) Comparison of the theoretical prediction of the backward transition process with numerical simulations. They agree with each other well, showing that the backward transition is universal, *i.e.*, independent of specific FDs.

or $\Theta_1^* = \arcsin(2/K)$ and $\Theta_2^* = \pi - \arcsin(2/K)$. Further linear analysis shows that the former steady solution is stable while the latter is unstable. For the amplitude equation, the steady solution satisfies the following equation:

$$(1 - r^{*2} - K\omega)r^* + K\omega r^*[1 + \cos\Theta^*]/2 = 0.$$
(9)

When $K \to \infty$, $\Theta^* \to 0$ according to eq. (8), thus $\cos \Theta^* \to 1$, and the steady solution of r should obey

$$1 - r^{*2} = 0, (10)$$

which gives $r^* = 1$ for the steady states.

Now, let us solve the order parameter R for the backward transition process. According to the definition,

$$R = \operatorname{Re}\left(\frac{1}{N}\sum_{j=1}^{N}r_{j}e^{i\eta_{j}}\right) = \frac{r^{*}}{2}(\cos\eta_{+}^{*} + \cos\eta_{-}^{*}). \quad (11)$$

Substituting eq. (5) and $r^* = 1$ into eq. (11), we get the analytical form of the order parameter as

$$R_1(K) = \frac{\sqrt{2}}{2} \sqrt{1 + \sqrt{1 - 4/K^2}},$$
 (12)

$$R_2(K) = \frac{\sqrt{2}}{2}\sqrt{1 - \sqrt{1 - 4/K^2}}.$$
 (13)

These two branches correspond to the stable and unstable solutions of eq. (8), respectively. Since R must be real, eqs. (12) and (13) imply that the (fully) coherent state in the system only exists for $K \geq 2$, *i.e.*, the OD state only exists when $K \geq 2$. If one continuously decreases K from ∞ , the stable and unstable OD states will gradually approach each other. Finally they collide and disappear via saddle-node bifurcation at $K_h^t = 2$ (the superscript t means theory), which just marks the first backward transition point in the system. Numerical results have verified that in all three cases of EOD, the backward transition occurs at $K_b \approx 2.02$, independently of the forms of specific FDs. In fig. 4, we compare the analytical curves of the order parameter with numerical results. It is shown that the backward transition processes with different FDs approximately follow a universal curve. The numerical and theoretical results are in quite good agreement with each other.

Conclusion. – In this paper, we reported the phenomenon of EOD in a system of coupled SL oscillators. By extensive numerical experiments, we classified three typical types of EOD, *i.e.*, OEOD, HEOD, and CEOD, corresponding to triangle, Lorentzian, and uniform FDs, respectively. In the case of OEOD, all oscillations suddenly cease at a critical point. In the case of HEOD, a few oscillators with large frequencies successively form two clusters first, then EOD suddenly occurs for the whole system. In the case of CEOD, there are two transitions towards OD, *i.e.*, first two giant clusters are formed in an explosive way; then they gradually absorb drifting oscillators until all oscillations are finally completely suppressed. We provided a self-consistent analysis, which enables us to predict a universal backward transition point, and prove that it is independent of the specific FDs. The theoretically results are supported by the numerical observations.

The present work demonstrated that, apart from the Kuramoto model, an explosive transition might occur in a certain dynamical system involving amplitudes. Therefore, it broadens our view of explosive phase transitions, and helps us to better understand the collective behaviors of networked systems. In our simulations, we have also observed that explosive OD will not occur in the present model if the network topology is scale free. This raises an important issue, *i.e.*, the interplay between network topology and dynamics in the process of OD/AD. In addition, could other coupling schemes also lead to explosive OD? Is it possible to observe explosive AD, and could such AD convert into OD [34]? Certainly, these questions deserve further investigation in the future.

* * *

This work is supported by the NSFC grants Nos. 11075056, 11375066, 11305062 and 11135001; the Innovation Program of Shanghai Municipal Education Commission grant No. 12ZZ043; and the Open Project Program of State Key Laboratory of Theoretical Physics, Institute of Theoretical Physics, Chinese Academy of Sciences, China (No. Y4KF151CJ1).

REFERENCES

- [1] ACHLIOPTAS D., D'SOUZA R. M. and SPENCER J., *Science*, **323** (2009) 1453.
- [2] ZIFF R. M., Phys. Rev. Lett., **103** (2009) 045701.
- [3] RADICCHI F. and FORTUNATO S., Phys. Rev. Lett., 103 (2009) 168701.
- [4] ARAÚJO N. A. M. and HERRMANN H. J., Phys. Rev. Lett., 105 (2010) 035701.

- [5] NAGLER J., LEVINA A. and TIMME M., Nat. Phys., 7 (2011) 265.
- [6] DA COSTA R. A., DOROGOVTSEV S. N., GOLTSEV A. V. and MENDES J. F. F., *Phys. Rev. Lett.*, **105** (2010) 255701.
- [7] RIORDAN O. and WARNKE L., Science, 333 (2011) 322.
- [8] GRASSBERGER P., CHRISTENSEN C., BIZHANI G., SON S.-W. and PACZUSKI M., Phys. Rev. Lett., 106 (2011) 225701.
- [9] CHO Y. S., HWANG S., HERRMANN H. J. and KAHNG B., Science, **339** (2013) 1185.
- [10] GÓMEZ-GARDEÑES J., GÓMEZ S., ARENAS A. and MORENO Y., Phys. Rev. Lett., 106 (2011) 128701.
- [11] LEYVA I. et al., Phys. Rev. Lett., 108 (2012) 168702.
- [12] PERON T. K. DM. and RODRIGUES F. A., Phys. Rev. E, 86 (2012) 056108.
- [13] LEYVA I. et al., Sci. Rep., 3 (2013) 1281.
- [14] LIU W., WU Y., XIAO J. and ZHAN M., EPL, 101 (2013) 38002.
- [15] JI P., PERON T. K. DM., MENCK P. J., RODRIGUES F. A. and KURTHS J., *Phys. Rev. Lett.*, **110** (2013) 218701.
- [16] ZHANG X., HU X., KURTHS J. and LIU Z., *Phys. Rev. E*, 88 (2013) 010802(R).
- [17] ZOU Y., PEREIRA T., SMALL M., LIU Z. and KURTHS J., *Phys. Rev. Lett.*, **112** (2014) 114102.
- [18] ZHANG X., ZOU Y., BOCCALETTI S. and LIU Z., Sci. Rep., 4 (2014) 5200.
- [19] SAXENA G., PRASAD A. and RAMASWAMY R., Phys. Rep., 521 (2012) 205.
- [20] KOSESKA A., VOLKOV E. and KURTHS J., Phys. Rep., 531 (2013) 173.
- [21] BAR-ELI K., Physica D, 14 (1985) 242.
- [22] MATTHEWS P. C., MIROLLO R. E. and STROGATZ S. H., *Physica D*, **52** (1991) 293.
- [23] ARONSON D. G., ERMENTROUT G. B. and KOPELL N., *Physica D*, **41** (1990) 403.
- [24] RAMANA REDDY D. V., SEN A. and JOHNSTON G. L., *Phys. Rev. Lett.*, **80** (1998) 5109.
- [25] ATAY F. M., Phys. Rev. Lett., 91 (2003) 094101.
- [26] LEE W. S., OTT E. and ANTONSEN T. M. jr., Chaos, 23 (2013) 033116.
- [27] RAMANA REDDY D. V., SEN A. and JOHNSTON G. L., *Phys. Rev. Lett.*, 85 (2000) 3381.
- [28] HERRERO R., FIGUERAS M., RIUS J., PI F. and ORRIOLS G., Phys. Rev. Lett., 84 (2000) 5312.
- [29] YANG J., Phys. Rev. E, 76 (2007) 016204.
- [30] ULLNER E., ZAIKIN A., VOLKOV E. I. and GARCÍA-OJALVO J., Phys. Rev. Lett., 99 (2007) 148103.
- [31] LIU W., WANG X., GUAN S. and LAI C.-H., New J. Phys., 11 (2009) 093016.
- [32] PRASAD A., DHAMALA M., ADHIKARI B. M. and RAMASWAMY R., *Phys. Rev. E*, 81 (2010) 027201.
- [33] KOSESKA A. and KURTHS J., Chaos, 20 (2010) 045111.
- [34] KOSESKA A., VOLKOV E. and KURTHS J., Phys. Rev. Lett., 111 (2013) 024103.